

Homotopy homomorphisms and the classifying space functor

R.M. Vogt

Abstract

We show that the classifying space functor $B : \mathcal{M}on \rightarrow \mathcal{Top}^*$ from the category of topological monoids to the category of based spaces is left adjoint to the Moore loop space functor $\Omega' : \mathcal{Top}^* \rightarrow \mathcal{M}on$ after we have localized $\mathcal{M}on$ with respect to all homomorphisms whose underlying maps are homotopy equivalences and $\mathcal{Top}^*$ with respect to all based maps which are (not necessarily based) homotopy equivalences. It is well-known that this localization of $\mathcal{Top}^*$ exists, and we show that the localization of $\mathcal{M}on$ is the category of monoids and homotopy classes of homotopy homomorphisms. To make this statement precise we have to modify the classical definition of a homotopy homomorphism, and we discuss the necessary changes. The adjunction is induced by an adjunction up to homotopy $B : \mathcal{H}\mathcal{M}on^w \rightleftarrows \mathcal{Top}^w : \Omega'$ between the category of well-pointed monoids and homotopy homomorphisms and the category of well-pointed spaces. This adjunction is shown to lift to diagrams. As a consequence, the well-known derived adjunction between the homotopy colimit and the constant diagram functor can also be seen to be induced by an adjunction up to homotopy before taking homotopy classes. As applications we among other things deduce a more algebraic version of the group completion theorem and show that the classifying space functor preserves homotopy colimits up to natural homotopy equivalences.

1 Introduction

Let $\mathcal{Top}$ denote the category of k -spaces, $\mathcal{Top}^*$ the category of based k -spaces, and $\mathcal{Top}^w$ the category of well-pointed k -spaces. Recall that a space X is a k -space if $A \subset X$ is closed iff $p^{-1}(A)$ is closed in C for each map $C \rightarrow X$

where C is a compact Hausdorff space, and that a space is called *well-pointed* if the inclusion of the base point is a closed cofibration.

Let $\mathcal{Mon}$ denote the category of topological monoids and continuous homomorphisms, and $\mathcal{Mon}^w$ and $\mathcal{CMon}$ the full subcategories of $\mathcal{Mon}$ of well-pointed respectively of commutative monoids. A monoid is canonically based by its unit.

We are interested in the relationship between Milgram's classifying space functor $B : \mathcal{Mon} \rightarrow \mathcal{Top}^*$ and the Moore loop space functor $\Omega' : \mathcal{Top}^* \rightarrow \mathcal{Mon}$ (for explicit definitions see Section 4).

If we restrict our attention to $\mathcal{CMon}$ we can easily establish a relationship between the classifying space functor and the usual loop space functor $\Omega : \mathcal{Top}^* \rightarrow \mathcal{Top}^*$: it is well-known that the classifying space BM of a commutative monoid is a commutative monoid [14], so that we have a functor $B : \mathcal{CMon} \rightarrow \mathcal{CMon}$. The usual loop space functor induces a functor $\Omega : \mathcal{CMon} \rightarrow \mathcal{CMon}$ by defining the multiplication in ΩM by point-wise multiplication in M . The category $\mathcal{CMon}$ is enriched over $\mathcal{Top}^*$ in an obvious way, and it is tensored and cotensored (for definitions see [6] or Section 3). The cotensor M^K of $M \in \mathcal{CMon}$ and $K \in \mathcal{Top}^*$ is the function space with point-wise multiplication. It is well-known that $B(M) \cong M \boxtimes S^1$, the tensor of M and S^1 . Since $- \boxtimes K$ is left adjoint to $(-)^K$ we obtain:

1.1 Proposition: The functors

$$B : \mathcal{CMon} \rightleftarrows \mathcal{CMon} : \Omega$$

form a $\mathcal{Top}^*$ -enriched adjoint pair.

In the non-commutative case there is no hope for a similar result. A candidate for a right adjoint of the classifying functor

$$B : \mathcal{Mon} \rightarrow \mathcal{Top}^*$$

is the Moore loop space functor

$$\Omega' : \mathcal{Top}^* \rightarrow \mathcal{Mon},$$

but Ω' does not preserve products. In fact, there is no product preserving functor

$$F : \mathcal{Top}^* \rightarrow \mathcal{Mon}$$

such that $F(X) \simeq \Omega(X)$ for all X [5, Prop. 6.1].

But Ω' does preserve products up to natural homotopy. So one might expect it to be a right adjoint of B up to homotopy or after a suitable localization of the categories. We will prove this in this paper.

1.2 Remark: This contrasts the situation in the simplicial category: The loop group functor $G : \mathcal{SSets} \rightarrow \mathcal{SGroups}$ from simplicial sets to simplicial groups is left adjoint to the simplicial classifying space functor $\overline{W} : \mathcal{SGroups} \rightarrow \mathcal{SSets}$ (e.g. see [11, Lemma V.5.3]).

We call a based map in $\mathcal{Top}^*$ a weak equivalence if it is a not necessarily based homotopy equivalence, and a homomorphism in $\mathcal{Mon}$ a weak equivalence if the underlying map of spaces is a weak equivalence in $\mathcal{Top}^*$. Let $\mathrm{Ho} \mathcal{Top}^*$ and $\mathrm{Ho} \mathcal{Mon}$ be the categories obtained from $\mathcal{Top}^*$ respectively $\mathcal{Mon}$ by formally inverting weak equivalences.

1.3 Theorem: The categories $\mathrm{Ho} \mathcal{Top}^*$ and $\mathrm{Ho} \mathcal{Mon}$ exist and the classifying space functor and the Moore loop space functor induce a derived adjoint pair

$$\mathrm{Ho} B : \mathrm{Ho} \mathcal{Mon} \rightleftarrows \mathrm{Ho} \mathcal{Top}^* : \mathrm{Ho} \Omega'$$

It is well-known that $\mathrm{Ho} \mathcal{Top}^*$ exists, but since it is not known that $\mathcal{Mon}$ admits a Quillen model category structure with our choice of weak equivalences the existence of $\mathrm{Ho} \mathcal{Mon}$ is a priori not clear. If we did choose the weak equivalences in $\mathcal{Top}^*$ to be weak homotopy equivalences we could avoid this problem, but we prefer our choice for two reasons: we obtain stronger results, namely genuine homotopy equivalences rather than weak ones, and our proofs have to be more geometric and therefore are of separate interest.

In the construction of $\mathrm{Ho} \mathcal{Mon}$ homotopy homomorphisms between monoids come into play. They were introduced by Sugawara in 1960 [21] and extensively studied by Fuchs in 1965 [8]. If we define a semigroup to be a topological space with a continuous associative multiplication, an inspection of the definition shows that a homotopy homomorphism $f : M \rightarrow N$ of monoids is nothing but a semigroup homomorphism $\overline{W}M \rightarrow N$ where $\overline{W}$ is a variant of the Boardman-Vogt W -construction [4] (not to be confused with the functor $\overline{W}$ of Remark 1.2). If $\mathcal{Sgp}$ denotes the category of semigroups and continuous homomorphisms then $\overline{W} : \mathcal{Sgp} \rightarrow \mathcal{Sgp}$ is a functor equipped with a natural transformation $\overline{\varepsilon} : \overline{W} \rightarrow \mathrm{Id}$. The Boardman-Vogt W -construction $W : \mathcal{Mon} \rightarrow \mathcal{Mon}$ and its associated natural transformation $\varepsilon : W \rightarrow \mathrm{Id}$ are obtained from $(\overline{W}, \overline{\varepsilon})$ by factoring out a unit relation. In particular, for any monoid M there is a natural projection $\varepsilon'(M) : \overline{W}M \rightarrow WM$ of semigroups such that $\varepsilon(M) \circ \varepsilon'(M) = \overline{\varepsilon}(M)$.

The lack of conditions for the unit is an indication that Sugawara's notion of a homotopy homomorphisms is not quite the correct one. So we define unitary homotopy homomorphism from M to N to be monoid homomorphisms $WM \rightarrow N$; those were studied in 1999 by Brinkmeier [3].

Composition of homotopy homomorphisms and their unitary versions is only associative up to homotopy. To obtain genuine categories of monoids and (unitary) homotopy homomorphisms we modify both notions: A homotopy homomorphisms from M to N will be a semigroup homomorphism $\overline{W}M \rightarrow \overline{W}N$ and a unitary one a monoid homomorphism $WM \rightarrow WN$. From a homotopy theoretical point of view this modification is not significant:

1.4 Proposition: If M, N are monoids and M is well-pointed and G, H are semigroups then the maps

$$\begin{aligned} \varepsilon(N)_* : \mathcal{M}on(WM, WN) &\rightarrow \mathcal{M}on(WM, N) \\ \bar{\varepsilon}(N)_* : \mathcal{S}gp(\overline{W}G, \overline{W}H) &\rightarrow \mathcal{S}gp(\overline{W}G, H) \end{aligned}$$

are homotopy equivalences.

It is well-known that $WM \rightarrow M$ has the flavor of a cofibrant replacement of M as known from model category theory provided M is well-pointed (e.g. see [2], [22]). So it is no surprise that the category of well-pointed monoids and homotopy classes of unitary homotopy homomorphisms is the localization of $\mathcal{M}on^w$ with respect to its weak equivalences. If we want to construct $\mathrm{Ho} \mathcal{M}on$ we have to laxen unitary homotopy homomorphisms to homotopy unitary homotopy homomorphisms and the corresponding statement holds. We will study these various notions of homotopy homomorphisms in Section 2 in detail.

If our categories lack the appropriate Quillen model structure, this is made up by their topological enrichment with nice properties. This topological enrichment allows us to prove stronger results. E.g. the restriction of Theorem 1.3 to the well-pointed case is the path-component version of the following result.

1.5 Theorem: Let $\mathcal{H}Mon^w$ be the category of well-pointed monoids and unitary homotopy homomorphisms. Then the classifying space functor and the Moore loop space functor induce an adjunction up to homotopy

$$\mathcal{H}Mon^w \rightleftarrows \mathcal{T}op^w.$$

In Section 3 we will introduce the necessary notions to make this precise. There we will also recall basic facts from enriched category theory and show that topologically enriched categories with a class of weak equivalences which admit a cofibrant replacement functor can be localized. We believe that these results are of separate interest.

In Section 4 we prove Theorem 1.5 and related results and hence Theorem 1.3. In Section 5 we draw some immediate consequences of Theorem 1.3 and of the intermediate steps in the proof of Theorem 1.5.

E.g. we obtain yet another but considerably shorter proof of a strong version of the James construction.

1.6 Definition: A *Dold space* is a topological space admitting a numerable cover $\{U_\gamma; \gamma \in \Gamma\}$ such that each inclusion $U_\gamma \subset X$ is nullhomotopic.

A space of the homotopy type of a *CW*-complex is a Dold space. For more details on Dold spaces see [17].

1.7 Proposition: (1) If X is a well-pointed space and JX is the based free topological monoid on X (the James construction), then $BJX \simeq \Sigma X$.
(2) If X is a well-pointed path-connected Dold space, then $JX \simeq \Omega \Sigma X$.

Part (2) was first proven in [7], shorter proofs can be found in [16] and [17].

We also obtain a new interpretation of the group completion theorem of a monoid without any additional assumptions on the multiplication.

1.8 Definition: A topological monoid is called *grouplike* if it admits a continuous homotopy inversion.

A standard example of a grouplike monoid is the Moore loop space $\Omega'X$ of a space X .

1.9 Theorem: Let M be a well-pointed topological monoid. Then there is a unitary homotopy homomorphism $\mu_M : M \rightarrow \Omega'BM$, natural up to homotopy, having the following universal property: Given any unitary homotopy homomorphism $f : M \rightarrow N$ into a grouplike monoid N there is a unitary homotopy homomorphism $\bar{f} : \Omega'BM \rightarrow N$, unique up to homotopy, such that $\bar{f} \circ \mu_M \simeq f$. (Here homotopy means homotopy in the category, i.e. homotopy through unitary homotopy homomorphisms.)

From the intermediate steps of the proof of Theorem 1.5 we obtain the following extension and strengthening of a theorem of Fuchs [8, Satz 7.7]

1.10 Proposition: (1) If M and N are well-pointed monoids and N is grouplike then

$$B : \mathcal{Mon}(WM, WN) \rightarrow \mathcal{Top}^*(BWM, BWN)$$

is a homotopy equivalence.

(2) If X is a well-pointed path-connected Dold space then $W\Omega' : \mathcal{Top}^w(X, Y) \rightarrow \mathcal{Mon}^w(W\Omega'X, W\Omega'Y)$ is a homotopy equivalence.

The reader may object that Fuchs considers homotopy homomorphisms while Proposition 1.10 addresses unitary homotopy homomorphisms. Since Fuchs only considers well-pointed grouplike monoids and all his spaces are of the homotopy type of CW -complexes the two notions are linked by

1.11 Proposition: Let M and N be well-pointed monoids and N be group-like. Then

$$(\varepsilon')^* : \mathcal{Mon}(WM, N) \rightarrow \mathcal{Sgp}(\overline{WM}, N)$$

is a homotopy equivalence,

Section 6 deals with diagrams in topologically enriched categories $\mathcal{M}$ with weak equivalences and a “good” cofibrant replacement functor. We first show that their localizations with respect to maps of diagrams which are object-wise weak equivalences exist. We then show that the well-known derived adjunction induced by the colimit functor and the constant diagram functor is the path-component version of an adjunction up to homotopy between the homotopy colimit functor and the constant diagram functor. We believe that this is of separate interest, too. We then show that the homotopy adjunction of Theorem 1.5 lifts to a homotopy adjunction between the corresponding categories of diagrams. In contrast to strict adjunctions this is a priori not clear, because the associated unit is natural only up to homotopy and hence does not lift to diagrams. We apply these result to prove

1.12 Theorem: The classifying space functor $B : \mathcal{Mon} \rightarrow \mathcal{Top}^*$ preserves homotopy colimits up to natural homotopy equivalence.

The path-component versions of most of our main results are more or less known if we restrict to grouplike monoids. The paper extends these results to general monoids and shows that they arise from stronger statements. Moreover, we show that a topological enrichment with good properties can make up for the non-existence of Quillen model structures.

2 Homotopy homomorphisms revisited

In 1960 Sugawara introduced the notion of a strongly homotopy multiplicative map between monoids, which we will call a homotopy homomorphisms

or h -morphism, for short [21].

2.1 Definition: A *homotopy homomorphism* or *h -morphism* $f : M \rightarrow N$ between two monoids is a sequence of maps

$$f_n : M^{n+1} \times I^n \longrightarrow N \quad n \in \mathbb{N}$$

such that $(x_i \in M, t_j \in I)$

$$\begin{aligned} f_n(x_0, t_1, x_1, t_2, \dots, t_n, x_n) \\ = \begin{cases} f_{n-1}(x_0, t_1, \dots, x_{i-1} \cdot x_i, \dots, t_n, x_n) & \text{if } t_i = 0 \\ f_{i-1}(x_0, t_1, \dots, x_{i-1}) \cdot f_{n-i}(x_i, t_{i+1}, \dots, x_n) & \text{if } t_i = 1 \end{cases} \end{aligned}$$

We call $f_0 : M \rightarrow N$ the *underlying map* of f .

If in addition $f_0(e_M) = e_N$ and

$$\begin{aligned} f_n(x_0, t_1, x_1, t_2, \dots, t_n, x_n) \\ = \begin{cases} f_{n-1}(x_1, t_2, \dots, x_n) & \text{if } x_0 = e_M \\ f_{n-1}(x_0, \dots, x_{i-1}, \max(t_i, t_{i+1}), x_{i+1}, \dots, x_n) & \text{if } x_i = e_M \\ f_{n-1}(x_0, t_1, \dots, x_{n-1}) & \text{if } x_n = e_M \end{cases} \end{aligned}$$

where $e_M \in M$ and $e_N \in N$ are the units, we call f a *unitary homotopy homomorphism* or *uh -morphism*, for short.

Since an h -morphism does not pay tribute to the unit it does not seem to be the right notion for maps between monoids. E.g. if we require f_0 to be a based map so that it preserves the unit we would like the path

$$\begin{array}{ccc} & f_1(x, t, y) & \\ \bullet & \xrightarrow{\quad\quad\quad} & \bullet \\ f_0(x \cdot y) & & f_0(x) \cdot f_0(y) \end{array}$$

to be the constant one, if x or y is the unit. Unitary h -morphisms have this property. Never-the-less, in the past one usually considered h -morphisms because the additional conditions for uh -morphisms make it harder to work with them.

We will later find it more convenient to work with homotopy unitary homotopy homomorphisms which preserve the unit only up to homotopy. We will introduce those at the end of this section.

The most extensive study of h -morphisms and their induced maps on classifying spaces was done by Fuchs [8], who constructed composites of h -morphisms, proved that composition is homotopy associative and stated that

an h -morphism $f : M \rightarrow N$ whose underlying map is a homotopy equivalence has a homotopy inverse h -morphisms $g : N \rightarrow M$. In fact, he constructed g_0, g_1 and the homotopies $g \circ f \simeq \text{id}$ and $f \circ g \simeq \text{id}$ in dimensions 0 and 1 in [8, p.205-p.208], but left the rest to the reader. He produced a complete proof in [9].

We handle these problems by interpreting homotopy homomorphisms as genuine homomorphisms of a “cofibrant” replacement of M .

By a *semigroup* we will mean a k -space with a continuous associative multiplication. Let $\mathcal{Sgp}$ denote the category of semigroups and continuous homomorphisms.

2.2 Constructions: We will construct continuous functors

$$\overline{W} : \mathcal{Sgp} \longrightarrow \mathcal{Sgp} \quad \text{and} \quad W : \mathcal{Mon} \longrightarrow \mathcal{Mon}$$

and natural transformations

$$\overline{\varepsilon} : \overline{W} \longrightarrow \text{Id} \quad \text{and} \quad \varepsilon : W \longrightarrow \text{Id}$$

as follows:

$$\overline{W}M = \left(\prod_{n=0}^{\infty} M^{n+1} \times I^n \right) / \sim$$

with the relation

$$(1) \quad (x_0, t_1, \dots, t_n, x_n) \sim (x_0, t_1, \dots, x_{i-1} \cdot x_i, \dots, t_n, x_n) \quad \text{if } t_i = 0$$

and WM is the quotient of $\overline{W}M$ by imposing the additional relations

$$(2) \quad \begin{aligned} & f_n(x_0, t_1, x_1, t_2, \dots, t_n, x_n) \\ & \sim \begin{cases} f_{n-1}(x_1, t_2, \dots, x_n) & \text{if } x_0 = e \\ f_{n-1}(x_0, \dots, x_{i-1}, \max(t_i, t_{i+1}), x_{i+1}, \dots, x_n) & \text{if } x_i = e \\ f_{n-1}(x_0, t_1, \dots, x_{n-1}) & \text{if } x_n = e \end{cases} \end{aligned}$$

The multiplications of $\overline{W}M$ and WM are given on representatives by

$$(x_0, t_1, \dots, x_k) \cdot (y_0, u_1, \dots, y_l) = (x_0, t_1, \dots, x_k, 1, y_0, u_1, \dots, y_l).$$

The natural transformations $\overline{\varepsilon}$ and ε are defined by

$$\overline{\varepsilon}(M), \varepsilon(M) : (x_0, t_1, \dots, x_k) \longmapsto x_0 \cdot x_1 \cdot \dots \cdot x_k.$$

Their underlying maps have natural sections

$$\overline{\iota}(M), \iota(M) : x \longmapsto (x)$$

which are not homomorphisms, and there is a homotopy over M

$$h_s : (x_0, t_1, x_1, \dots, x_n) \longmapsto (x_0, s \cdot t_1, \dots, s \cdot t_n, x_n)$$

from $\bar{\iota}(M) \circ \bar{\varepsilon}(M)$ respectively $\iota(M) \circ \varepsilon(M)$ to the identity. In particular, $\bar{\varepsilon}(M)$ and $\varepsilon(M)$ are shrinkable as maps.

If M is a monoid the projection

$$\varepsilon'(M) : \bar{W}M \rightarrow WM$$

is a homomorphism of semigroups satisfying

$$\bar{\varepsilon}(M) = \varepsilon(M) \circ \varepsilon'(M) \quad \text{and} \quad \varepsilon'(M) \circ \bar{\iota}(M) = \iota(M).$$

By inspection we see

2.3 Observation: (1) h -morphisms $(f_n) : M \rightarrow N$ correspond bijectively to homomorphisms $\bar{f} : \bar{W}M \rightarrow N$ of semigroups, and $f_0 = \bar{f} \circ \bar{\iota}(M)$

(2) uh -morphisms $(f_n) : M \rightarrow N$ correspond bijectively to homomorphisms $f : WM \rightarrow N$ of monoids, and $f_0 = f \circ \iota(M)$

2.4 Observation: Algebraically, $\bar{W}M$ is a free semigroup and WM is a free monoid. The indecomposables are precisely those elements which have a representative $(x_0, t_1, x_1, \dots, x_n)$ where no t_i equals 1.

2.5 The formal relation between $\bar{W}$ and W : The forgetful functor $i : \mathcal{Mon} \rightarrow \mathcal{Sgp}$ has a left adjoint

$$(-)_+ : \mathcal{Sgp} \rightarrow \mathcal{Mon}, \quad G \mapsto G_+,$$

where $G_+ = G \sqcup \{*\}$ with $*$ as unit. It follows from the definitions that the diagram

$$\begin{array}{ccc} \mathcal{Sgp} & \xrightarrow{\bar{W}} & \mathcal{Sgp} \\ (-)_+ \downarrow & & \downarrow (-)_+ \\ \mathcal{Mon} & \xrightarrow{W} & \mathcal{Mon} \end{array}$$

commutes up to natural isomorphisms in $\mathcal{Mon}$.

Both constructions have a universal property, which is a consequence of the following result. We give $\mathcal{Top}^*(X, Y)$ and $\mathcal{Top}(X, Y)$ the k -function space topology, obtained by turning the space of all maps from X to Y with the compact-open topology into a k -space. We give $\mathcal{Mon}(M, N)$ and $\mathcal{Sgp}(M, N)$ the subspace topologies of the corresponding function spaces in $\mathcal{Top}^*$ respectively $\mathcal{Top}$.

2.6 Definition: We call a homomorphism $f : M \rightarrow N$ in $\mathcal{Mon}$ or $\mathcal{Sgp}$ a *weak equivalence* if its underlying map of spaces is a homotopy equivalence in $\mathcal{Top}$. (Recall that a weak equivalence in $\mathcal{Mon}$ is a homotopy equivalence of underlying spaces in $\mathcal{Top}^*$ if M and N are well-pointed.)

2.7 Proposition: (1) Let M be a well-pointed monoid and $p : X \rightarrow Y$ a homomorphism of monoids. Let

$$p_* : \mathcal{Mon}(WM, X) \longrightarrow \mathcal{Mon}(WM, Y)$$

be the induced map. If p is a fibration of underlying spaces, so is p_* . If p is a weak equivalence, p_* is a homotopy equivalence.

(2) The same holds for $\overline{W}$ and an arbitrary object M in the category $\mathcal{Sgp}$.

Proof Let $p : X \rightarrow Y$ be a weak equivalence. By the HELP-Lemma [23] in $\mathcal{Top}$ with the Strøm model structure [19] we have to show: Given a diagram of spaces

(A)

$$\begin{array}{ccc} A & \xrightarrow{\bar{f}_A} & \mathcal{Mon}(WM, X) \\ \downarrow i & & \downarrow p_* \\ B & \xrightarrow{\bar{g}} & \mathcal{Mon}(WM, Y) \end{array}$$

which commutes up to a homotopy $h_{A,t} : \bar{g} \circ i \simeq p_* \circ \bar{f}_A$, where i is a closed cofibration, there are extensions $\bar{f} : B \rightarrow \mathcal{Mon}(WM, X)$ of $\bar{f}_A$ and $\bar{h}_t : B \rightarrow \mathcal{Mon}(WM, Y)$ of $\bar{h}_{A,t}$ such that $\bar{h}_t : \bar{g} \simeq p_* \circ \bar{f}$.

Passing to adjoints we obtain a diagram

$$\begin{array}{ccc} WM \times A & \xrightarrow{f_A} & X \\ \downarrow id \times i & & \downarrow p \\ WM \times B & \xrightarrow{g} & Y \end{array}$$

commuting up to a homotopy $h_{A,t}$, such that each $f_a = f_A|_{WM \times \{a\}}$, each $g_b = g|_{WM \times \{b\}}$, and each $h_{a,t} = h_{A,t}|_{WM \times \{a\}}$ is a homomorphism. We have to construct extensions $f : WM \times B \rightarrow X$ and $h_t : WM \times B \rightarrow Y$ of f_A and $h_{A,t}$ such that $h_t : g \simeq p \circ f$ and each $h_{b,t}$ and f_b , $b \in B$ is a homomorphism.

We filter $WM \times B$ by closed subspaces $F_n \times B$, where F_n is the submonoid of WM generated by all elements having a representative $(x_0, t_1, \dots, t_k, x_k)$

with $k \leq n$. We put $F_{-1} = \{e\}$. Then f and h_t are uniquely determined on $F_{-1} \times B$.

Now suppose that f and h_t have been defined on $F_{n-1} \times B$. An element $(x_0, t_1, \dots, t_n, x_n)$ represents an element in F_{n-1} iff one of the following conditions holds

- some $x_i = e$ (relation 2.2.2)
- some $t_i = 0$ (relation 2.2.1)
- some $t_i = 1$ (it represents a product in F_{n-1}).

If $DM^{n+1} \subset M^{n+1}$ denotes the subspace of points with some coordinate e , then f and h_t are already defined on $(DM^{n+1} \times I^n \cup M^{n+1} \times \partial I^n) \times B \cup M^{n+1} \times I^n \times A$. The elements in $(M^{n+1} \times I^n) \setminus (DM^{n+1} \times I^n \cup M^{n+1} \times \partial I^n)$ represent indecomposables of filtration n , but not of lower filtration. Consider the diagram

(B)

$$\begin{array}{ccc} (DM^{n+1} \times I^n \cup M^{n+1} \times \partial I^n) \times B \cup M^{n+1} \times I^n \times A & \xrightarrow{j} & M^{n+1} \times I^n \times B \\ \downarrow f & & \downarrow g \\ X & \xrightarrow{p} & Y \end{array}$$

(in abuse of notation we use g for the composite $M^{n+1} \times I^n \times B \rightarrow WM \times B \rightarrow Y$). Diagram (B) commutes up to the homotopy h_t and we need an extension of f and h_t to $M^{n+1} \times I^n \times B$. These extensions exist by the HELP-Lemma, because our assumptions ensure that j is a closed cofibration. So we have defined f and h_t for indecomposable generators $(x_0, t_1, \dots, t_n, x_n)$ of F_n . We extend these maps to $F_n \times B$ by the conditions that each f_b and $h_{b,t}$, $b \in B$ be a homomorphism using Observation 2.4.

Now suppose that p is a fibration. By [19, Thm. 8] we need to consider a commutative diagram (A), where i is a closed cofibration and a homotopy equivalence, and we have to find an extension $\bar{f} : B \rightarrow \mathcal{Mon}(WM, X)$ of $\bar{f}_A$ such that $\bar{g} = p_* \circ \bar{f}$. We proceed as above. In the inductive step we have a commutative diagram (B). Since i is a closed cofibration and a homotopy equivalence so is j by the pushout-product theorem for cofibrations. Hence the required extension $f : M^{n+1} \times I^n \times B \rightarrow X$ exists by [19, Thm. 8].

Part (2) is proved in the same way starting with $F_{-1}M = \emptyset$. $\square$

As an immediate consequence we obtain the

2.8 Lifting Theorem: Given homomorphisms of monoids

$$\begin{array}{ccc} & & X \\ & & \downarrow p \\ WM & \xrightarrow{f} & Y \end{array}$$

such that p is a weak equivalence and M is well-pointed, then there exists a homomorphism $g : WM \rightarrow X$, unique up to homotopy in $\mathcal{Mon}$ (i.e. a homotopy through homomorphisms) such that $f \simeq p \circ g$ in $\mathcal{Mon}$.

If, in addition, the underlying map of p is a fibration there is a homomorphism $g : WM \rightarrow X$, unique up to homotopy in $\mathcal{Mon}$, such that $f = p \circ g$.

(2) For $\overline{W}$ the analogous results hold in the category $\mathcal{Sgp}$.

2.9 By Proposition 2.7 the second one of the maps

$$\begin{aligned} \varepsilon(N)_* : \mathcal{Mon}(WM, WN) &\rightarrow \mathcal{Mon}(WM, N) \\ \overline{\varepsilon}(N)_* : \mathcal{Sgp}(\overline{W}M, \overline{W}N) &\rightarrow \mathcal{Sgp}(\overline{W}M, N) \end{aligned}$$

is a homotopy equivalence, and the first one is a homotopy equivalence if M is well-pointed.

To guarantee the well-pointedness condition we introduce the whiskering functor.

2.10 The whiskering costruction: We define a functor

$$V^t : \mathcal{Top}^* \rightarrow \mathcal{Top}^w$$

by $V^t(X, x_0) = (X \sqcup I)/(x_0 \sim 1)$ and choose $0 \in I$ as base-point of X_I . Then $V^t X$ is wellpointed, and the natural map $q(X) : V^t X \rightarrow X$ mapping I to x_0 is a homotopy equivalence. If X is wellpointed, $q(X)$ is a based homotopy equivalence.

This functor lifts to a functor

$$V : \mathcal{Mon} \rightarrow \mathcal{Mon}^w$$

defined by $V(M) = V^t(M)$ with x_0 replaced by e_M with the multiplication

$$x \cdot y = \begin{cases} x \cdot y \in M & \text{if } x, y \in M \\ x & \text{if } x \in M, y \in I \\ y & \text{if } y \in M, x \in I \\ \max(x, y) & \text{if } x, y \in I \end{cases}$$

Since $0 \in I$ is the unit of VM the monoid VM is well-pointed. The natural map $q(M) : VM \rightarrow M$ is a weak equivalence in $\mathcal{Mon}$.

A homomorphism $f : WVM \rightarrow N$ can be considered a homotopy unitary homotopy homomorphism with underlying map $f_0 = f \circ \iota(M) \circ \bar{q}(M) : M \rightarrow VM \rightarrow WVM \rightarrow N$, where $\bar{q}(M) : M \rightarrow VM$ is the canonical map. We note that f_0 preserves the unit e_M only up to homotopy.

By 2.9 the following **change of our notations of homotopy homomorphisms** is insignificant from a homotopy theoretic point of view:

2.11 Definition: From now on a *homotopy unitary homotopy homomorphism*, *huh*-morphism for short, from M to N is a homomorphism $f : WVM \rightarrow WVN$. Its *underlying map* is $\varepsilon(N) \circ f \circ \iota(M) \circ \bar{q}(M)$.

A *unitary homotopy homomorphism*, *uh*-morphism for short, from M to N is a homomorphism $f : WM \rightarrow WN$. Its *underlying map* is $\varepsilon(N) \circ f \circ \iota(M)$.

A *homotopy homomorphism*, *h*-morphism for short, from the semigroup M to the semigroup N is a homomorphism $f : \overline{WM} \rightarrow \overline{WN}$. Its *underlying map* is $\bar{\varepsilon}(N) \circ f \circ \bar{\iota}(M)$.

This solves the problem of composition, and from 2.7 we obtain

2.12 Proposition: If $f : WM \rightarrow WN$ is a *uh*-morphism from M to N whose underlying map is a homotopy equivalence, and M and N are well-pointed, then f is a homotopy equivalence in the category $\mathcal{Mon}$.

If $f : WVM \rightarrow WVN$ is a *huh*-morphism from M to N , whose underlying map is a homotopy equivalence, then f is a homotopy equivalence in the category $\mathcal{Mon}$.

The analogous statement in $\mathcal{Sgp}$ holds for homomorphisms $\overline{WM} \rightarrow \overline{WN}$.

3 Categorical prerequisites and localizations

The functors $WV : \mathcal{Mon} \rightarrow \mathcal{Mon}$ and $\overline{W} : \mathcal{Sgp} \rightarrow \mathcal{Sgp}$ resemble cofibrant replacement functors as known from Quillen model category theory. Unfortunately, there is no known model category structure on $\mathcal{Mon}$ with our choice of weak equivalences. This draw-back is made up by the topological enrichment of our categories as we will see in this section.

Our categories are enriched over $\mathcal{Top}^*$ or $\mathcal{Top}$. So we have a natural notion of homotopy. Moreover, they are *tensor*ed and *cotensor*ed. Recall that a $\mathcal{Top}^*$ -enriched category $\mathcal{M}$ is *tensor*ed and *cotensor*ed (over $\mathcal{Top}^*$) if there are functors

$$\begin{aligned} \mathcal{Top}^* \times \mathcal{M} &\rightarrow \mathcal{M}, & (X, M) &\mapsto X \boxtimes M \\ (\mathcal{Top}^*)^{\text{op}} \times \mathcal{M} &\rightarrow \mathcal{M}, & (X, M) &\mapsto M^X \end{aligned}$$

and natural homeomorphism

$$\mathcal{M}(X \boxtimes M, N) \cong \mathcal{Top}^*(X, \mathcal{M}(M, N)) \cong \mathcal{M}(M, N^X).$$

These properties imply that for based spaces X and Y and objects $M \in \mathcal{M}$ there are natural isomorphisms

$$(X \wedge Y) \boxtimes M \cong X \boxtimes (Y \boxtimes M).$$

The definition in the $\mathcal{Top}$ -enriched case is accordingly. To distinguish between the based and the non-based case we denote the tensor over $\mathcal{Top}$ by $X \otimes M$. The natural isomorphism in non-based case reads

$$(X \times Y) \otimes M \cong X \otimes (Y \otimes M).$$

Forgetting base points turns a $\mathcal{Top}^*$ -enriched category $\mathcal{M}$ into a $\mathcal{Top}$ -enriched one. If $\mathcal{M}$ is tensored over $\mathcal{Top}^*$ it is also tensored over $\mathcal{Top}$: we define

$$X \otimes M = X_+ \boxtimes M$$

where $X_+ = X \sqcup \{*\}$ with the additional point as base point.

3.1 Example: $\mathcal{Mon}$ is $\mathcal{Top}^*$ -enriched, tensored and cotensored [15, Prop. 2.10]. The cotensor M^X is the k -function space with pointwise multiplication, $X \boxtimes M$ is more complicated: as a set, it is a free product of copies M , one copy for each $x \in X$ different from the base point. By the same argument as in [15] the category $\mathcal{Sgp}$ is $\mathcal{Top}$ -enriched and tensored and cotensored over $\mathcal{Top}$.

If $\otimes_{\mathcal{Sgp}}$ denotes the tensor in $\mathcal{Sgp}$ and $\otimes$ the one over $\mathcal{Top}$ in $\mathcal{Mon}$, then the universal properties of the tensor and of the adjunction of 2.5 imply that there is a natural isomorphism

$$(K \otimes_{\mathcal{Sgp}} G)_+ \cong K \otimes (G_+)$$

in $\mathcal{Mon}$ for semigroups G .

If $\mathcal{M}$ is tensored over $\mathcal{Top}$ it has a canonical cylinder functor $M \mapsto I \otimes M$. The associated homotopy notion coincides with the natural one. We denote the quotient category of $\mathcal{M}$ obtained by passing to homotopy classes by $\pi\mathcal{M}$.

3.2 Definition: Let $\mathcal{M}$ be a category and $\mathcal{W}$ a class of morphisms in $\mathcal{M}$, which we will call *weak equivalences*. The *localization* of $\mathcal{M}$ with respect to $\mathcal{W}$ is a category $\mathcal{M}[\mathcal{W}^{-1}]$ with $\text{ob } \mathcal{M}[\mathcal{W}^{-1}] = \text{ob } \mathcal{M}$ and a functor $\gamma : \mathcal{M} \rightarrow$

$\mathcal{M}[\mathcal{W}^{-1}]$ such that

- (1) γ is the identity on objects
- (2) $\gamma(f)$ is an isomorphism for all $f \in \mathcal{W}$
- (3) if $F : \mathcal{M} \rightarrow \mathcal{D}$ is a functor such that $F(f)$ is an isomorphism for all $f \in \mathcal{W}$ then there exists a unique functor $\overline{F} : \mathcal{M}[\mathcal{W}^{-1}] \rightarrow \mathcal{D}$ such that $F = \overline{F} \circ \gamma$.

3.3 Proposition: Let $\mathcal{M}$ be a $\mathcal{T}op$ -enriched tensored category and $\mathcal{W}$ a class of morphisms in $\mathcal{M}$ such that

- (1) $\mathcal{W}$ contains all homotopy equivalences,
- (2) there is a functor $Q : \mathcal{M} \rightarrow \mathcal{M}$ and a natural transformation $\varepsilon : Q \rightarrow \text{Id}$ or a natural transformation $\eta : \text{Id} \rightarrow Q$ taking values in $\mathcal{W}$ such that Qf is a homotopy equivalence for each $f \in \mathcal{W}$.

Then $\mathcal{M}[\mathcal{W}^{-1}]$ exists. More precisely, $\mathcal{M}[\mathcal{W}^{-1}] = \pi\mathcal{HC}$, where $\text{ob } \mathcal{HC} = \text{ob } \mathcal{M}$ and $\mathcal{HM}(M_1, M_2) = \mathcal{M}(QM_1, QM_2)$, and $\gamma : \mathcal{M} \rightarrow \mathcal{M}[\mathcal{W}^{-1}]$ maps a morphism f to the homotopy class of Qf .

Proof The proof is the same as in the case of a Quillen model category (e.g. see [10, Thm 8.3.5]). Observe that γ maps weak equivalences to isomorphisms by assumption (2), and that a functor $F : \mathcal{D} \rightarrow \mathcal{D}$ which maps weak equivalences to isomorphisms maps homotopic maps to the same map. The latter holds because the bottom and top inclusions $i_0, i_1 : X \rightarrow I \otimes X$ into the cylinder are homotopy equivalences with a common homotopy inverse. $\square$

3.4 Remark: For Proposition 3.3 we do not need that the tensor $X \otimes M$ exists for all topological spaces: it suffices that $I \otimes M$ and $I^2 \otimes M$ exist so that the above argument works.

3.5 Notation: Following the standard convention we denote $\mathcal{M}[\mathcal{W}^{-1}]$ by $\text{Ho } \mathcal{M}$ if the class $\mathcal{W}$ has been specified.

A pair $(Q, \varepsilon : Q \rightarrow \text{Id})$ respectively $(Q, \eta : \text{Id} \rightarrow Q)$ satisfying the requirements of 3.3 will be called a *cofibrant* respectively *fibrant replacement functor*. The $\mathcal{T}op$ -enriched categories $\mathcal{M}$ considered in this paper will have a continuous cofibrant replacement functor, and we call the category $\mathcal{HM}$ the *category of Q -morphisms* associated with $\mathcal{M}$.

3.6 Definition: A functor $Q : \mathcal{M} \rightarrow \mathcal{M}$ together with a natural transformation $\varepsilon : Q \rightarrow \text{Id}$ is called a *strong cofibrant replacement functor* if each $\varepsilon(M) : Q(M) \rightarrow M$ is a weak equivalence and $p_* : \mathcal{M}(QA, B) \rightarrow \mathcal{M}(QA, C)$ is a homotopy equivalence whenever $p : B \rightarrow C$ is a weak equivalence.

Clearly, a strong cofibrant replacement functor is a cofibrant replacement functor.

3.7 Examples:

1. Let $\mathcal{W} \subset \mathcal{Mon}$ be the class of weak equivalences in the sense of 2.6. Then $WV : \mathcal{Mon} \rightarrow \mathcal{Mon}$ together with $WVM \xrightarrow{\varepsilon(VM)} VM \xrightarrow{q(M)} M$ is a strong cofibrant replacement functor by 2.12, and the Q -morphisms are the huh -morphisms.
2. Let $\mathcal{W} \subset \mathcal{Mon}^w$ be again the class of weak equivalences. Then $W : \mathcal{Mon}^w \rightarrow \mathcal{Mon}^w$ together with $\varepsilon : W \rightarrow \text{Id}$ is a strong cofibrant replacement functor by 2.12, and the Q -morphisms are the uh -morphisms.
3. Let $\mathcal{W} \subset \mathcal{Sgp}$ be the class of weak equivalences. Then $\overline{W} : \mathcal{Sgp} \rightarrow \mathcal{Sgp}$ together with $\overline{\varepsilon} : \overline{W} \rightarrow \text{Id}$ is a strong cofibrant replacement functor by 2.12, and the Q -morphisms are the h -morphisms.
4. Let $\mathcal{W} \subset \mathcal{Top}^*$ be the class of based maps which are (not necessarily based) homotopy equivalences. Then $V^t : \mathcal{Top}^* \rightarrow \mathcal{Top}^*$ together with $q : V^t \rightarrow \text{Id}$ is a strong cofibrant replacement functor by the lemma below, the proof of which we leave as an exercise.
5. Let $\mathcal{W} \subset \mathcal{Top}^w$ be the class of homotopy equivalences. Then $\text{Id} : \mathcal{Top}^w \rightarrow \mathcal{Top}^w$ is a strong cofibrant replacement functor and each map is a Q -morphism.

3.8 Lemma: Let A be a well-pointed space and $p : X \rightarrow Y$ a map in $\mathcal{Top}^*$ which is a not necessarily based homotopy equivalence. Then

$$p_* : \mathcal{Top}^*(A, X) \rightarrow \mathcal{Top}^*(A, Y)$$

is a homotopy equivalence in $\mathcal{Top}$.

3.9 Proposition: The localizations of the categories of 3.7 with respect to their weak equivalences exist.

This follows from 3.3 and 3.4, if we can show that the tensors $I \otimes M$ and $I^2 \otimes M$ exist in $\mathcal{Mon}^w$, which holds by [15, Prop. 7.8].

3.10 Definition: Let $\mathcal{M}$ be a category and $\mathcal{W}$ a class of morphisms in $\mathcal{M}$ such that $\mathcal{M}[\mathcal{W}^{-1}]$ exists. Let $F : \mathcal{M} \rightarrow \mathcal{D}$ be a functor. A functor $\mathbf{L}F : \mathcal{M}[\mathcal{W}^{-1}] \rightarrow \mathcal{D}$ together with a natural transformation $\tau : \mathbf{L}F \circ \gamma \rightarrow F$

is called *left derived functor* of F , if given any functor $T : \mathcal{M}[\mathcal{W}^{-1}] \rightarrow \mathcal{D}$ and natural transformation $\sigma : T \circ \gamma \rightarrow F$, there is a unique natural transformation $\rho : T \rightarrow \mathbf{L}F$ such that $\sigma = \tau \circ (\rho * \gamma)$.

Dually, a functor $\mathbf{R}F : \mathcal{M}[\mathcal{W}^{-1}] \rightarrow \mathcal{D}$ together with a natural transformation $\mu : F \rightarrow \mathbf{R}F \circ \gamma$ is called *right derived functor* of F , if given any functor $G : \mathcal{M}[\mathcal{W}^{-1}] \rightarrow \mathcal{D}$ and natural transformation $\nu : F \rightarrow G \circ \gamma$, there is a unique natural transformation $\xi : \mathbf{R}F \rightarrow G$ such that $(\xi * \gamma) \circ \mu$.

3.11 Remark: (1) A left or right derived functor is unique up to natural isomorphism if it exists.

(2) If $F : \mathcal{M} \rightarrow \mathcal{D}$ maps weak equivalences to isomorphisms, then the induced functor $\overline{F} : \mathcal{M}[\mathcal{W}^{-1}] \rightarrow \mathcal{D}$ is the right and left derived functor of F .

3.12 Proposition: Let $\mathcal{M}$ be as in Proposition 3.3, and let $F : \mathcal{M} \rightarrow \mathcal{B}$ be a functor which maps homotopy equivalences to isomorphisms. Then $\mathbf{L}(\pi \circ F) : \mathcal{M}[\mathcal{W}^{-1}] \rightarrow \mathcal{B}$ exists if $\mathcal{M}$ has a cofibrant replacement functor, and $\mathbf{R}F : \mathcal{M}[\mathcal{W}^{-1}] \rightarrow \mathcal{B}$ exists if $\mathcal{M}$ has a fibrant replacement functor. In both cases the derived functor is induced by $F \circ Q : \mathcal{M} \rightarrow \mathcal{B}$.

Proof The proof is the same as in the case of a model category (e.g. see [10, 8.4.]). $\square$

3.13 Let $F : \mathcal{M} \rightarrow \mathcal{B}$ be a functor between $\mathcal{Top}$ -enriched categories admitting cofibrant replacement functors $Q_{\mathcal{M}} : \mathcal{M} \rightarrow \mathcal{M}$ and $Q_{\mathcal{B}} : \mathcal{B} \rightarrow \mathcal{B}$. Proposition 3.12 motivates the introduction of the functor

$$F^{\mathcal{H}} : \mathcal{H}\mathcal{M} \rightarrow \mathcal{H}\mathcal{B}$$

defined on objects by $F^{\mathcal{H}}(X) = F(Q_{\mathcal{M}}X)$ and on morphisms by

$$F^{\mathcal{H}} : \mathcal{M}(Q_{\mathcal{M}}X, Q_{\mathcal{M}}Y) \xrightarrow{Q_{\mathcal{B}} \circ F} \mathcal{B}(Q_{\mathcal{B}}FQ_{\mathcal{M}}X, Q_{\mathcal{B}}FQ_{\mathcal{M}}Y).$$

If F preserves homotopy equivalences, e.g. if F is continuous, and $\pi_{\mathcal{B}} : \mathcal{B} \rightarrow \pi\mathcal{B}$ is the canonical functor, then $\pi_{\mathcal{B}} \circ F^{\mathcal{H}}$ induces the left derived functor

$$\mathrm{Ho} F : \mathrm{Ho} \mathcal{M} \rightarrow \mathrm{Ho} \mathcal{B}$$

of $\pi_{\mathcal{B}} \circ F$. Following model category terminology, we call $\mathrm{Ho} F$ the *total left derived functor* of F .

One of the objectives of this paper is to show that the classifying space functor and the Moore loop space functor induce an adjoint derived pair (see

Theorem 4.6 below). This is the path-component version of the more general result (Theorem 4.5 below) that

$$B^{\mathcal{H}} : \mathcal{H}Mon \rightleftarrows \mathcal{T}op^* : \Omega^{\mathcal{H}}$$

are a homotopically adjoint pair. To make this last statement precise we need some preparations.

3.14 Definition: Let $\mathcal{A}$ and $\mathcal{B}$ be topologically enriched categories. A functor $F : \mathcal{A} \rightarrow \mathcal{B}$ is called *continuous* if

$$F : \mathcal{A}(A, B) \longrightarrow \mathcal{B}(FA, FB)$$

is continuous for all A and B in $\mathcal{A}$.

If $F, G : \mathcal{A} \rightarrow \mathcal{B}$ are continuous functors, a collection of morphisms $\{\alpha(A) : FA \rightarrow GA; A \in \text{ob } \mathcal{A}\}$ is called a *natural transformation up to homotopy* if the diagram

$$\begin{array}{ccc} \mathcal{A}(A, B) & \xrightarrow{F} & \mathcal{B}(FA, FB) \\ \downarrow G & & \downarrow \alpha(B)_* \\ \mathcal{B}(GA, GB) & \xrightarrow{\alpha(A)^*} & \mathcal{B}(FA, GB) \end{array}$$

is homotopy commutative.

A pair of continuous functors

$$F : \mathcal{A} \rightleftarrows \mathcal{B} : G$$

is called a *homotopy adjoint pair* if there is a natural transformation up to homotopy

$$\alpha(A, X) : \mathcal{B}(FA, X) \rightarrow \mathcal{A}(A, GX)$$

such that each $\alpha(A, X)$ is a homotopy equivalence.

Observe that we have chosen a strong form of a natural transformation $\alpha : F \rightarrow G$ up to homotopy: for each morphism $f : A \rightarrow B$ in $\mathcal{A}$ we have a square

$$\begin{array}{ccc} FA & \xrightarrow{\alpha(A)} & GA \\ Ff \downarrow & & \downarrow Gf \\ FB & \xrightarrow{\alpha(B)} & GB \end{array}$$

commuting up to a homotopy $H(f)$ which is continuous in f .

The proofs of the following two lemmas are easy exercises.

3.15 Lemma: Let $S, T, U : \mathcal{A} \rightarrow \mathcal{B}$ be continuous functors of topologically enriched categories.

- (1) Each natural transformation $\alpha : S \rightarrow T$ is a natural transformations up to homotopy.
- (2) If $\varepsilon : S \rightarrow T$ and $\eta : T \rightarrow U$ are natural transformations up to homotopy, then $\eta \circ \varepsilon : S \rightarrow U$ is one.
- (3) Let $\varepsilon : S \rightarrow T$ be a natural transformation up to homotopy such that each $\varepsilon(A)$ is a homotopy equivalence. Choose a homotopy inverse $\eta(A)$ of $\varepsilon(A)$ for each A in $\mathcal{A}$. Then the $\eta(A)$ form a natural transformation $\eta : T \rightarrow S$ up to homotopy. $\square$

3.16 Lemma: Let $S, T, U, V : \mathcal{A} \rightarrow \mathcal{B}$ be continuous functors of topologically enriched categories, and let $\varepsilon : S \rightarrow T$ and $\eta : U \rightarrow V$ be natural transformations up to homotopy.

- (1) Let $F, G : \mathcal{A}^{\text{op}} \times \mathcal{A} \rightarrow \mathcal{Top}$ be defined by $F(A, B) = \mathcal{A}(A, B)$ and $G(A, B) = \mathcal{B}(TA, TB)$. Then

$$\tau(A, B) : \mathcal{A}(A, B) \xrightarrow{T} \mathcal{B}(TA, TB)$$

is a natural transformation from F to G .

- (2) Let $F, G : \mathcal{A}^{\text{op}} \times \mathcal{A} \rightarrow \mathcal{Top}$ be defined by $F(A, B) = \mathcal{B}(VA, SB)$ and $G(A, B) = \mathcal{B}(UA, TB)$. Then

$$\alpha(A, B) : \mathcal{B}(VA, SB) \xrightarrow{\varepsilon(B)_* \circ \eta(A)^*} \mathcal{B}(UA, TB)$$

is a natural transformation from F to G up to homotopy. $\square$

3.17 Proposition: Let $F : \mathcal{A} \rightleftarrows \mathcal{B} : G$ be a pair of continuous functors of topologically enriched categories. Suppose there are natural transformations up to homotopy

$$\mu(A) : A \rightarrow GF(A) \quad \text{and} \quad \eta(X) : FG(X) \rightarrow X$$

such that

$$G(\eta(X)) \circ \mu(GX) \simeq \text{id}_{GX} \quad \text{and} \quad \eta(FA) \circ (F(\mu(A))) \simeq \text{id}_{FA}.$$

Then F and G are a homotopy adjoint pair. (We call $\mu : \text{Id} \rightarrow GF$ the *homotopy unit* and $\eta : FG \rightarrow \text{Id}$ the *homotopy counit* of the resulting homotopy adjunction.)

Proof We define

$$\alpha(A, X) : \mathcal{B}(FA, X) \xrightarrow{G} \mathcal{A}(GFA, GX) \xrightarrow{\mu(A)^*} \mathcal{A}(A, GX)$$

and

$$\beta(A, X) : \mathcal{A}(A, GX) \xrightarrow{F} \mathcal{B}(FA, FGX) \xrightarrow{\eta(X)_*} \mathcal{B}(FA, X).$$

By 3.16 both are natural transformations up to homotopy. The following diagram shows that $\beta(A, X) \circ \alpha(A, X) \simeq \text{id}$.

$$\begin{array}{ccccc}
& & \mathcal{A}(GFA, GX) & \xrightarrow{\mu(A)^*} & \mathcal{A}(A, GX) \\
& \nearrow G & \downarrow F & \text{II} & \downarrow F \\
\mathcal{B}(FA, X) & \text{I} & \mathcal{B}(FGFA, FGX) & \xrightarrow{(F\mu(A))^*} & \mathcal{B}(FA, FGX) \\
& \searrow \eta(FA)^* & \downarrow \eta(X)_* & \text{III} & \downarrow \eta(X)_* \\
& & \mathcal{B}(FGFA, X) & \xrightarrow{(F\mu(A))^*} & \mathcal{B}(FA, X)
\end{array}$$

The squares II and III commute and square I commutes up to homotopy, and $(F\mu(A))^* \circ \eta(FA)^* \simeq \text{id}$ by assumption.

The proof that $\alpha(A, X) \circ \beta(A, X) \simeq \text{id}$ is dual. $\square$

3.18 Definition: A homotopy adjunction $F : \mathcal{A} \rightleftarrows \mathcal{B} : G$ is called *natural* if there is a natural homotopy equivalence

$$\beta(A, X) : \mathcal{A}(A, GX) \rightarrow \mathcal{B}(FA, X)$$

and *conatural* if there is a natural homotopy equivalence

$$\alpha(A, X) : \mathcal{B}(FA, X) \rightarrow \mathcal{A}(A, GX)$$

(because in this case there is a natural homotopy unit respectively a natural homotopy counit).

4 The classifying space and the Moore loop space functor

4.1 The 2-sided bar construction: Let $\mathcal{C}$ be a small topologically enriched category, X a $\mathcal{C}^{op}$ -diagram and Y a $\mathcal{C}$ -diagram in $\mathcal{T}op$. We define a

simplicial space $B_\bullet(X, \mathcal{C}, Y)$ by

$$\begin{aligned} B_0(X, \mathcal{C}, Y) &= \coprod_{A \in \mathcal{C}} X(A) \times Y(A) \\ B_n(X, \mathcal{C}, Y) &= \coprod_{A, B \in \mathcal{C}} X(B) \times \mathcal{C}_n(A, B) \times Y(A) \quad \text{for } n > 0, \end{aligned}$$

where $\mathcal{C}_n(A, B)$ is the space of all composable n -tuples of morphisms $(f_1, \dots, f_n)$ such that $\text{source}(f_n) = A$ and $\text{target}(f_1) = B$, with boundary and degeneracy maps given by

$$\begin{aligned} d^i(x, f_1, \dots, f_n, y) &= (X(f_1)(x), f_2, \dots, f_n, y) & i = 0 \\ d^i(x, f_1, \dots, f_n, y) &= (x, f_1, \dots, f_i \circ f_{i+1}, \dots, f_n, y) & 0 < i < n \\ d^i(x, f_1, \dots, f_n, y) &= (x, f_1, \dots, f_{n-1}, Y(f_n)(y)) & i = n \\ s^i(x, f_1, \dots, f_n, y) &= (x, f_1, \dots, f_i, \text{id}, f_{i+1}, \dots, f_n, y) & 0 \leq i \leq n \end{aligned}$$

Let $B(X, \mathcal{C}, Y) = |B_\bullet(X, \mathcal{C}, Y)|$ be its topological realization.

We consider a topological monoid as a topologically enriches category with one object and define the *classifying space functor*

$$B : \mathcal{Mon} \longrightarrow \mathcal{Top}^*$$

by $BM = B(*, M, *)$. Since BM is well-pointed if M is, the classifying space functor is a functor of pairs

$$B : (\mathcal{Mon}, \mathcal{Mon}^w) \rightarrow (\mathcal{Top}^*, \mathcal{Top}^w).$$

4.2 We will also work with the variant

$$\tilde{B} : \mathcal{Mon} \longrightarrow \mathcal{Top}^*$$

where the topological realizations of $B_\bullet(*, M, *)$ is replaced by the fat realization which disregards degeneracies. Since the fat realization does not make use of identities the functor $\tilde{B}$ extends to $\mathcal{Sgp}$; moreover, $\tilde{B}G$ is well-pointed for any semigroup G so that

$$\tilde{B} : \mathcal{Sgp} \rightarrow \mathcal{Top}^w.$$

By construction, there is a natural homeomorphism $\tilde{B}(G) \cong B(G_+)$ for semi-groups G , and the diagram

$$\begin{array}{ccc} \tilde{B}(M) & \xrightarrow{\cong} & B(M_+) \\ & \searrow p(M) & \swarrow B(\kappa(M)) \\ & B(M) & \end{array}$$

commutes for monoids M , where $\kappa : M_+ \rightarrow M$ is the counit of the adjunction 2.5 and $p : \tilde{B} \rightarrow B$ is the natural projection.

It is well-known that $p(M) : \tilde{B}(M) \rightarrow B(M)$ and hence $B(\kappa(M)) : B(M_+) \rightarrow B(M)$ are homotopy equivalences if M is well-pointed.

4.3 The Moore path and loop space: Let X be a (not necessarily based) space. The *Moore path space* of X is the subspace $\text{Path}(X) \subset X^{\mathbb{R}_+} \times \mathbb{R}_+$ consisting of all pairs (w, r) such that $w(t) = w(r)$ for all $t \geq r$. We call r the *length* of w and denote it by $r = l(w)$.

For two paths (w_1, r_1) and (w_2, r_2) with $(w_1, r_1)(r_1) = (w_2, r_2)(0)$ we define *path addition* by

$$(w_1, r_1) + (w_2, r_2) = (w, r_1 + r_2)$$

with

$$w(t) = \begin{cases} w_1(t), & 0 \leq t \leq r_1, \\ w_2(t - r_1) & r_1 \leq t. \end{cases}$$

If $(X, *)$ is a based space, the *Moore loop space* $\Omega'(X) \subset \text{Path}(X)$ is the subspace of all pairs (w, r) with $(w, r)(r) = (w, r)(0) = *$. Path addition defines a monoid structure on $\Omega'X$ with $(c, 0)$ as unit, where $c : \mathbb{R}_+ \rightarrow X$ is the constant map to $*$. The usual loop space ΩX is embedded in $\Omega'(X)$ as a deformation retract.

It follows from [7, (11.3)] that $\Omega'(X)$ is well-pointed if X is. Hence Ω' defines a functor of pairs

$$\Omega' : (\mathcal{Top}^*, \mathcal{Top}^w) \rightarrow (\mathcal{Mon}, \mathcal{Mon}^w).$$

Following 3.13 we have pairs of continuous functors

$$B^{\mathcal{H}} : \mathcal{HMon} \rightleftarrows \mathcal{HTop}^* : \Omega'^{\mathcal{H}}$$

and

$$B^{w\mathcal{H}} : \mathcal{HMon}^w \rightleftarrows \mathcal{HTop}^w = \mathcal{Top}^w : \Omega'^{w\mathcal{H}}.$$

We shall prove

4.4 Theorem: The functors

$$B^{w\mathcal{H}} : \mathcal{HMon}^w \rightleftarrows \mathcal{Top}^w : \Omega'^{w\mathcal{H}}$$

are a conatural homotopically adjoint pair: There is a continuous natural map

$$\lambda(WM, X) : \mathcal{Mon}(WM, W\Omega'X) \longrightarrow \mathcal{Top}^*(BWM, X)$$

which is a homotopy equivalence.

As an immediate consequence we obtain

4.5 Theorem: The functors

$$B^{\mathcal{H}} : \mathcal{H}Mon \rightleftarrows \mathcal{H}Top^* : \Omega'^{\mathcal{H}}$$

are a conatural homotopically adjoint pair: There is a continuous natural map

$$\lambda(WVM, V^tX) : Mon(WVM, WV\Omega'V^tX) \longrightarrow Top^*(V^tBWVM, V^tX)$$

which is a homotopy equivalence.

Proof Replacing M by VM and X by V^tX in Proposition 4.4 we obtain a natural homotopy equivalence

$$Mon(WVM, WV\Omega'V^tX) \xrightarrow{\sim} Top^*(BWVM, V^tX).$$

Since $BWVM$ is well-pointed the natural map $q(BWVM) : V^tBWVM \rightarrow BWVM$ is a based homotopy equivalence inducing a natural homotopy equivalence

$$q(BWVM)^* : Top^*(BWVM, V^tX) \rightarrow Top^*(V^tBWVM, V^tX).$$

□

Passing to homotopy classes we obtain

4.6 Theorem: The functors

$$Ho B : Ho Mon \rightleftarrows Ho Top^* : Ho \Omega'$$

are an adjoint pair. Moreover, $Ho B$ is the left derived of $\gamma_{Top^*} \circ B$ and $Ho \Omega'$ the left derived of $\gamma_{Mon} \circ \Omega'$.

Proof This follows from our explicit description of the localizations and the derived functors in Section 3. □

The rest of this Section is devoted to the proof of Theorem 4.4. By 3.17 it suffices to construct a homotopy unit $\mu : Id_{\mathcal{H}Mon^w} \rightarrow \Omega'^{w\mathcal{H}}B^{w\mathcal{H}}$ and a homotopy counit $\eta : B^{w\mathcal{H}}\Omega'^{w\mathcal{H}} \rightarrow Id_{\mathcal{H}Top^w}$. Then $\lambda(WM, X)$ is the composite

$$Mon(WM, W\Omega'X) \xrightarrow{B} Top^*(BW M, BW\Omega'X) \xrightarrow{\eta(X)^*} Top^*(BW M, X).$$

4.7 This means, we have to construct continuous homomorphisms

$$\mu(WM) : WM \longrightarrow W\Omega'BMW$$

which constitute a natural transformation up to homotopy with respect to homomorphisms $WM \rightarrow WN$, and a natural transformation

$$\eta(X) : BW\Omega'X \longrightarrow X,$$

such that

(1) $W\Omega'\eta(X) \circ \mu(W\Omega'X) \simeq \text{id}_{W\Omega'X}$ in $\mathcal{Mon}^w$ and

(2) $\eta(BWM) \circ B\mu(WM) \simeq \text{id}_{BWM}$ in $\mathcal{Top}^w$.

(For λ to be a natural transformation we need η to be a natural transformation.)

4.8 The homotopy counit: Let X be a based space and let

$$\Delta^n = \{(t_0, \dots, t_n) \in \mathbb{R}^{n+1}; \sum_{i=0}^n t_i = 1, t_i \geq 0 \text{ for all } i\}$$

denote the standard n -simplex. The *evaluation map*

$$\text{ev}(X) : B\Omega'X = \left(\coprod_{n \geq 0} (\Omega'X)^n \times \Delta^n \right) / \sim \longrightarrow X$$

is defined by

$$\text{ev}(X)((w_1, \dots, w_n)(t_0, \dots, t_n)) = (w_1 + \dots + w_n) \left(\sum_{i=1}^n t_i \cdot \sum_{j=1}^i l(w_j) \right)$$

where $l(w_j)$ is the length of w_j .

The homotopy counit η is the natural map

$$\eta(X) : BW\Omega'X \xrightarrow{B\varepsilon(\Omega'X)} B\Omega'X \xrightarrow{\text{ev}(X)} X.$$

4.9 The homotopy unit: For a monoid M let EM denote the 2-sided bar construction $B(M, M, *)$. Then

$$z \cdot (x_0, x_1, \dots, x_n) = (z \cdot x_0, x_1, \dots, x_n)$$

defines a left M -action on the simplicial space $B_\bullet(M, M, *)$ and hence on EM .

Let $P(EM)$ denote the space of Moore paths in EM starting at the base-point (e) in the 0-skeleton M of EM . The endpoint projection

$$P(EM) \longrightarrow EM$$

is known to be a fibration. Moreover, it is a homotopy equivalence because $P(EM)$ and EM are contractible. Let $P(EM, M)$ be the pullback

$$\begin{array}{ccc} P(EM, M) & \longrightarrow & P(EM) \\ \downarrow \pi(M) & & \downarrow \\ M & \xrightarrow{i} & EM \end{array}$$

where i is the inclusion of the 0-skeleton, i.e. $P(EM, M)$ is the space of Moore paths in EM starting at (e) and ending in M . Then $\pi(M)$ is a fibration and a homotopy equivalence. We define a monoid structure $\oplus$ in $P(EM, M)$ by

$$w_1 \oplus w_2 = w_1 + x \cdot w_2$$

where $+$ is the usual path addition, $x \in M$ is the endpoint of w_1 , and $x \cdot w_2$ is the path $t \mapsto x \cdot w_2(t)$. Then $\pi(M) : P(EM, M) \rightarrow M$ is a homomorphism and hence a weak equivalence of monoids.

Factoring out the operation of M on EM we obtain a projection

$$EM \rightarrow BM$$

inducing a homomorphism

$$\rho'(M) : (P(EM, M), \oplus) \longrightarrow (\Omega'BM, +).$$

Since we do not know whether $(P(EM, M))$ is well-pointed or not we apply the whiskering process to it and obtain a homomorphism

$$\rho(M) : V(P(EM, M), \oplus) \xrightarrow{q((P(EM, M)))} (P(EM, M), \oplus) \xrightarrow{\rho'(M)} (\Omega'BM, +).$$

The homomorphism $\sigma(M) : WVP(EM, M) \rightarrow M$ defined by

$$\begin{array}{ccccc} WVP(EM, M) & \xrightarrow{\varepsilon(VP(EM, M))} & VP(EM, M) & \xrightarrow{q(P(EM, M))} & P(EM, M) \\ & \searrow \sigma(M) & & & \swarrow \pi(M) \\ & & M & & \end{array}$$

is a weak equivalence. All these constructions are functorial in M and the maps between them are natural in M . We apply them to WM rather than to M ; in particular $\sigma(WM)$ is a homotopy equivalence in $\mathcal{Mon}^w$.

We choose a homotopy inverse of $\sigma(WM)$ in $\mathcal{Mon}^w$

$$\nu(WM) : WM \longrightarrow WVP(EM, WM),$$

which is a natural transformation up to homotopy with respect to homomorphisms $WM \rightarrow WN$ by Lemma 3.15.

We define our homotopy unit by

$$\mu(WM) : WM \xrightarrow{\nu(WM)} WVP(EM, WM) \xrightarrow{W\rho(WM)} W\Omega'BM,$$

which is a natural transformation up to homotopy by Lemma 3.15.

Our verification of the conditions 4.7 depends on an explicit description of an h -morphism $M \rightarrow \Omega'BM$ defined by a natural homomorphism

$$\zeta'(M) : \overline{W}(M) \longrightarrow \Omega'BM$$

and the interplay of $\overline{W}(M)$ and WM .

We define $\zeta'(M)$ as a composite of homomorphisms

$$\overline{W}(M) \xrightarrow{\zeta(M)} P(EM, M) \xrightarrow{\rho'(M)} \Omega'BM$$

The homomorphism $\zeta(M)$ maps the element represented by $(x_0, t_1, \dots, x_n)$ to the path

$$v_0 + v_1 + \dots + v_n$$

of length $t_1 + \dots + t_n + 1$ in the simplex $(e, x_0, x_1, \dots, x_n) \times \Delta^{n+1} \subset EM$, where

$$v_k(s) = (e, x_0, \dots, x_n) \times (u_0, \dots, u_{n+1}) \quad \text{and} \quad l(v_k) = t_{k+1}$$


with

$$u_r = \begin{cases} (1-s) \cdot t_r \cdot \prod_{j=r+1}^k (1-t_j) & r \leq k \\ s & r = k+1 \\ 0 & r \geq k+2 \end{cases}$$

and the convention that $t_{n+1} = 1$.

Observe that $+$ is the usual path addition of Moore paths in EM and not the monoid structure of $P(EM, M)$.


Example: $(x_0, t_1, x_1, t_2, x_2)$ is mapped to the path $v_0 + v_1 + v_2$ of length $t_1 + t_2 + 1$ given by



4.10 By construction, $\pi(M) \circ \zeta(M) = \bar{\varepsilon}(M)$. In particular, $\zeta(M) : \overline{WM} \rightarrow P(EM, M)$ is a weak equivalence of semigroups.

4.11 Remark: We will show below that $\rho(M) : P(EM, M) \rightarrow \Omega'BM$ is a weak equivalence if M is grouplike, so that $\rho(M) \circ \zeta(M)$ is an h -morphism which is a weak equivalence if M is grouplike. It is well-known that such an h -morphism exists, but to our knowledge there is no explicit description in the literature.

4.12 Consider the following diagram



where $\sigma_1(WM) = q(P(EM, M)) \circ \varepsilon(VP(EM, M))$ and $f^+ : G_+ \rightarrow M$ is the adjoint of the homomorphism $f : G \rightarrow M$ from a semigroup into a monoid.

By definition of $\nu(WM)$ and $\sigma(WM)$ the left lower triangle commutes up to homotopy in $\mathcal{Mon}^w$ and the right lower triangle is commutative. Since

$$\pi(WM) \circ \zeta(WM) = \bar{\varepsilon}(WM) \simeq \pi(WM) \circ \sigma_1(WM) \circ \nu(WM) \circ \bar{\varepsilon}(WM)$$

Proposition 2.7 implies that

$$\sigma_1(WM) \circ \nu(WM) \circ \bar{\varepsilon}(WM) \simeq \zeta(WM) \quad \text{in } \mathcal{Sgp}$$

which in turn is equivalent to the saying that square I commutes up to homotopy in $\mathcal{Mon}$.

We are now in the position to prove

4.13 Proposition: $\eta(BWM) \circ B\mu(WM) \simeq \text{id}_{BWM}$ in $\mathcal{Top}^w$.

This result is a fairly easy consequence of

4.14 Lemma: The diagram

$$\begin{array}{ccccccc} \tilde{B}\overline{W}M & \xrightarrow{\tilde{B}\bar{\varepsilon}(M)} & & \tilde{B}M & & & \\ \tilde{B}\zeta(M) \downarrow & & & \downarrow p(M) & & & \\ \tilde{B}P(EM, M) & \xrightarrow{\tilde{B}\rho'(M)} & \tilde{B}\Omega'BM & \xrightarrow{p(\Omega'BM)} & B\Omega'BM & \xrightarrow{\text{ev}(BM)} & BM \end{array}$$

commutes up to homotopy.

Proof Let $f = \text{ev}(BM) \circ p(\Omega'BM) \circ \tilde{B}\rho'(M) \circ \tilde{B}\zeta$ and let $g = p(M) \circ \tilde{B}\bar{\varepsilon}(M)$. Let $z = (z_1, \dots, z_n)$ be an element in $(\overline{W}M)^n$, so that $z \times \Delta^n$ is an n -simplex in $\tilde{B}\overline{W}M$. If $z_j = (x_{j0}, t_{j1}, \dots, x_{jr_j})$, then f maps $z \times \Delta^n$ to the image of the path $\rho'(M) \circ \zeta(z_1) + \dots + \rho'(M) \circ \zeta(z_n)$ which lies in the simplex

$$\sigma = \sigma(z) = (x_{10}, \dots, x_{1r_1}, \dots, x_{n0}, \dots, x_{nr_n}) \times \Delta^{r_1 + \dots + r_n + n}$$

in BM , while g maps $z \times \Delta^n$ identically (modulo possible degenerations) onto the simplex

$$\tau = \tau(z) = (x_{10} \cdot \dots \cdot x_{1r_1}, \dots, x_{n0} \cdot \dots \cdot x_{nr_n}) \times \Delta^n$$

in BM , which is a face of σ . So $f|_{z \times \Delta^n}$ is homotopic to $g|_{z \times \Delta^n}$ by a linear homotopy. We call a homotopy from f to g *admissible* if it maps $z \times \Delta^n$ to $\sigma(z)$ throughout the homotopy.

We are going to construct an admissible homotopy $H : \widetilde{B\overline{W}M} \times I \rightarrow BM$ from f to g by induction on the canonical filtration $(\widetilde{B\overline{W}M})^{(n)}$ of $\widetilde{B\overline{W}M}$.

$(\widetilde{B\overline{W}M})^{(0)}$ is a point, which is mapped by f and g to the base-point. Now suppose that we have constructed an admissible homotopy

$$H : (\widetilde{B\overline{W}M})^{(n-1)} \times I \rightarrow BM.$$

Let $z \times \Delta^n$ be an n -simplex in $\widetilde{B\overline{W}M}$ as above. We define

$$q(z) = q(z_1, \dots, z_n) = r_1 + \dots + r_n \in \mathbb{N}$$

and we extend H over $(\widetilde{B\overline{W}M})^{(n)} \times I$ by induction on q .

If $q = 0$, then $z = (z_1, \dots, z_n)$ with $z_j = (x_{j0})$ for $j = 1, \dots, n$ and $\sigma(z) = \tau(z) = (x_{10}, \dots, x_{n0}) \times \Delta^n$. Hence the space of all n -simplices $z \in (\widetilde{B\overline{W}M})^{(n)}$ with $q(z) = 0$ is M^n . By induction, we have to find a homotopy

$$h : M^n \times \Delta^n \times I \rightarrow M^n \times \Delta^n$$

over M^n which is already determined on $M^n \times \partial(\Delta^n \times I)$. If b_n denotes the barycenter of Δ^n we map $((x_1, \dots, x_n), b_n, \frac{1}{2})$ to $((x_1, \dots, x_n), b_n)$ and cone off.

If $q > 0$ we have q coordinates $t_{jk} \in I$ in z . So the space of all elements z with $q(z) = q$ is the union of spaces of the form $M^{n+q} \times I^q$ which may intersect on their lower faces $M^{n+q} \times LI^q$ due to the relations, where $LI^q = \{(t_1, \dots, t_q) \in I^q; \text{ some } t_i = 0\}$. So possible intersections are of lower filtration. We have to find a map

$$h : M^{n+q} \times I^q \times \Delta^n \times I \rightarrow M^{n+q} \times \Delta^{n+q}$$

over M^{n+q} which is already defined on

$$M^{n+q} \times (LI^q \times \Delta^n \times I \cup I^q \times \partial(\Delta^n \times I)).$$

Since LI^q is a strong deformation retract of I^q , the inclusion

$$LI^q \times \Delta^n \times I \cup I^q \times \partial(\Delta^n \times I) \subset I^q \times \Delta^n \times I$$

is an inclusion of a strong deformation retract. Hence h exists. $\square$

Proof of Proposition 4.13: Since M is well-pointed, the projection $p(M) : \widetilde{B}M \rightarrow BM$ is a homotopy equivalence. If $h : X \rightarrow Y$ is a weak equivalence of semigroups, then $\widetilde{B}h : \widetilde{B}X \rightarrow \widetilde{B}Y$ is a based homotopy equivalence. Hence it suffices to show that

$$\eta(BWM) \circ B\mu(WM) \circ p(WM) \circ \widetilde{B}\overline{\varepsilon}(WM) \simeq p(WM) \circ \widetilde{B}\overline{\varepsilon}(WM).$$

Using Diagram 4.12, Lemma 4.14 with M replaced by WM , and the naturality of ε and p we have

$$\begin{aligned}
& \eta(BWM) \circ B\mu(WM) \circ p(WM) \circ \tilde{B}\bar{\varepsilon}(WM) \\
&= \text{ev}(BWM) \circ B\varepsilon(\Omega' BWM) \circ BW\rho(WM) \circ B\nu(WM) \circ p(WM) \circ \tilde{B}\bar{\varepsilon}(WM) \\
&= \text{ev}(BWM) \circ p(\Omega' BWM) \circ \tilde{B}\rho(WM) \circ \tilde{B}\varepsilon(VP(EWM, WM)) \circ \tilde{B}\nu(WM) \circ \tilde{B}\bar{\varepsilon}(WM) \\
&\simeq \text{ev}(BWM) \circ p(\Omega' BWM) \circ \tilde{B}\rho'(WM) \circ \tilde{B}\zeta(WM) \\
&\simeq p(WM) \circ \tilde{B}\bar{\varepsilon}(WM). \quad \square
\end{aligned}$$

The proof of the first part of 4.7 needs some preparation. Let $\mathcal{J}$ denote the category of ordered sets $[n] = \{0 < 1 < \dots < n\}$ and order preserving injections, and let $\mathcal{JTop}_0$ denote the category of all diagrams

$$X_\bullet : \mathcal{J}^{\text{op}} \rightarrow \mathcal{Top}, \quad [n] \mapsto X_n$$

such that X_0 is a single point, i.e. an object in $\mathcal{JTop}_0$ is a reduced simplicial space without degeneracies. Of lately, such an object is called a *reduced semisimplicial space*. The usual fat topological realization functor

$$\mathcal{JTop}_0 \rightarrow \mathcal{Top}^*, \quad X_\bullet \mapsto \|X_\bullet\|$$

has a right adjoint, the *reduced singular functor*

$$\text{Sing}_\bullet^0 : \mathcal{Top}^* \rightarrow \mathcal{JTop}_0, \quad \text{Sing}_n^0(Y) = \mathcal{Top}((\Delta^n, \Delta_0^n), (Y, *))$$

where Δ_0^n is the 0-skeleton of Δ^n . The unit of this adjunction

$$\tau_\bullet(X_\bullet) : X_\bullet \rightarrow \text{Sing}_\bullet^0 \|X_\bullet\|$$

sends $x \in X_n$ to the singular simplex

$$\Delta^n \xrightarrow{i_x} \coprod_k X_k \times \Delta^k \rightarrow \|X_\bullet\|$$

where i_x is the inclusion of the simplex $\{x\} \times \Delta^n$. The counit

$$\widehat{\text{ev}}(Y) : \| \text{Sing}_\bullet^0(Y) \| \rightarrow Y$$

is induced by the evaluation maps $\text{Sing}_n^0(Y) \times \Delta^n \rightarrow Y$. The formula defining our evaluation map of 4.8 defines a natural semisimplicial map

$$\alpha_\bullet(Y) : N_\bullet \Omega' Y \rightarrow \text{Sing}_\bullet^0$$

where $N_\bullet \Omega'Y$ is the semisimplicial nerve of $\Omega'Y$. Let $v_0, \dots, v_n$ denote the vertices of Δ^n and let $L_n \subset \Delta^n$ denote the union of the 1-simplexes $[v_{i-1}, v_i]$, $i = 1, \dots, n$. Then L_n is a strong deformation retract of Δ^n . The composite

$$(\Omega'Y)^n = N_n(\Omega'Y) \xrightarrow{\alpha_n(Y)} \text{Sing}_n^0(Y) \xrightarrow{r} \text{Top}((L_n, \Delta_0), (Y, *)) = (\Omega Y)^n,$$

where r is the restriction to L_n , is the map normalizing the loop lengths to 1. In particular, $\alpha_n(Y)$ is a homotopy equivalence inducing a homotopy equivalence $\|\alpha_\bullet(Y)\|$. Moreover, the diagram

4.15

$$\begin{array}{ccc} \tilde{B}\Omega'Y & \xrightarrow{p(\Omega'Y)} & B\Omega'Y \\ \|\alpha_\bullet(Y)\| \downarrow & & \downarrow \text{ev}(Y) \\ \|\text{Sing}_\bullet^0(Y)\| & \xrightarrow{\widehat{\text{ev}}(Y)} & Y \end{array}$$

commutes.

4.16 Proposition: (1) If M is a grouplike well-pointed monoid, then $\rho(M)$ and hence $\mu(WM)$ are weak equivalences.

(2) If Y is a well-pointed path-connected Dold space (see Definition 1.6), then $\text{ev}(Y) : B\Omega'Y \rightarrow Y$ and hence $\eta(Y)$ is a based homotopy equivalence.

(3) If Y is a well-pointed space, then $\Omega' \text{ev}(Y) : \Omega' B\Omega'Y \rightarrow \Omega'Y$ is a weak equivalence. Hence so is $\Omega'\eta(Y)$.

(4) If M is a well-pointed monoid, then $B\mu(WM) : BW M \rightarrow BW \Omega' BW M$ is a homotopy equivalence.

Proof (1) The diagram

$$\begin{array}{ccccc} M & \xrightarrow{\tau_1(N_\bullet M)} & \Omega \tilde{B}M & \xrightarrow{\Omega p(M)} & \Omega BM \\ \zeta \circ \bar{i}(M) \downarrow & & & & \downarrow i(BM) \\ P(EM, M) & \xrightarrow{\rho(M)} & & & \Omega' BM \end{array}$$

commutes. Here $i(X) : \Omega X \rightarrow \Omega' X$ is the inclusion and $\bar{i}(M) : M \rightarrow \overline{W}M$ the section (see 2.2). It is well known that $\tau_1(N_\bullet M)$ is a homotopy equivalence if M is grouplike (e.g. see [18]). Since $p(M)$, $i(BM)$, and $\zeta \circ \bar{i}(M)$ are homotopy equivalences in Top , so is $\rho(M)$.

(2) In the commutative diagram 4.15 the map $p(\Omega'Y)$ is a homotopy equivalence because $\Omega'Y$ is well-pointed and $\widehat{\text{ev}}(Y)$ is a homotopy equivalence by [17, Prop. 5.6].

(3) Consider the following commutative diagram in $\mathcal{JTop}_0$

$$\begin{array}{ccc}
N_{\bullet}\Omega'Y & \xrightarrow{\tau_{\bullet}(N_{\bullet}\Omega'Y)} & \text{Sing}_{\bullet}^0 \widetilde{B}\Omega'Y \\
\alpha_{\bullet}(Y) \downarrow & & \downarrow \text{Sing}_{\bullet}^0 \|\alpha_{\bullet}(Y)\| \\
\text{Sing}_{\bullet}^0 Y & \xrightarrow{\tau_{\bullet}(\text{Sing}_{\bullet}^0 Y)} & \text{Sing}_{\bullet}^0 \parallel \text{Sing}_{\bullet}^0 Y \parallel \\
& \searrow \text{id} & \downarrow \text{Sing}_{\bullet}^0 \widehat{\text{ev}}(Y) \\
& & \text{Sing}_{\bullet}^0 Y
\end{array}$$

Restricting this diagram to degree 0 we obtain a commutative diagram of spaces

$$\begin{array}{ccccc}
\Omega'Y & \xrightarrow{\tau_1(N_{\bullet}\Omega'Y)} & \Omega \widetilde{B}\Omega'Y & \xrightarrow{\Omega p(\Omega'Y)} & \Omega B\Omega'Y \\
\alpha_1 \downarrow & & \downarrow \Omega \|\alpha_{\bullet}(Y)\| & & \downarrow \Omega \text{ev}(Y) \\
\Omega Y & \xrightarrow{\tau_1(\text{Sing}_{\bullet}^0 Y)} & \Omega \parallel \text{Sing}_{\bullet}^0 Y \parallel & \xrightarrow{\Omega \widehat{\text{ev}}(Y)} & \Omega Y.
\end{array}$$

Since $\Omega'Y$ is grouplike, $\tau_1(N_{\bullet}\Omega'Y)$ is a homotopy equivalence. Since Y and hence $\Omega'Y$ is well-pointed, $\Omega p(\Omega'Y)$ is a homotopy equivalence. Since α_1 and $\|\alpha_{\bullet}(Y)\|$ are homotopy equivalences, $\tau_1(\text{Sing}_{\bullet}^0 Y)$ is one. Hence so is $\Omega \widehat{\text{ev}}(Y)$ and hence also $\Omega \text{ev}(Y)$, which implies the result.

(4) Since BWM is a well-pointed path-connected Dold space by [17, Cor. 5.2] the statement follows from Part (2) and Proposition 4.13. $\square$

4.17 Proposition: $W\Omega'\eta(X) \circ \mu(W\Omega'X) \simeq \text{id}_{W\Omega'X}$ in $\mathcal{Mon}$.

Proof It follows from Proposition 4.13 and the homotopy naturality of μ and η that the following diagram commutes up to homotopy.

$$\begin{array}{ccccc}
W\Omega'X & \xrightarrow{\mu(W\Omega'X)} & W\Omega'BW\Omega'X & \xrightarrow{W\Omega'\eta(X)} & W\Omega'X \\
\mu(W\Omega'X) \downarrow & & \downarrow \mu(W\Omega'BW\Omega'X) & & \downarrow \mu(W\Omega'X) \\
W\Omega'BW\Omega'X & \xrightarrow{W\Omega'B\mu(W\Omega'X)} & W\Omega'BW\Omega'BW\Omega'X & \xrightarrow{W\Omega'BW\Omega'\eta(X)} & W\Omega'BW\Omega'X \\
& \searrow \text{id} & \downarrow W\Omega'\eta(BW\Omega'X) & & \downarrow W\Omega'\eta(X) \\
& & W\Omega'BW\Omega'X & \xrightarrow{W\Omega'\eta(X)} & W\Omega'X
\end{array}$$

We obtain

$$W\Omega'\eta(X) \circ \mu(W\Omega'X) \circ W\Omega'\eta(X) \circ \mu(W\Omega'X) \simeq W\Omega'\eta(X) \circ \mu(W\Omega'X).$$

Since $\Omega'X$ is grouplike $\mu(W\Omega'X)$ and $W\Omega'\eta(X)$ are weak equivalences by Proposition 4.16. By Proposition 2.12 both homomorphisms have homotopy inverses in $\mathcal{Mon}$ so that

$$W\Omega'\eta(X) \circ \mu(W\Omega'X) \simeq \text{id}_{W\Omega'X}$$

in $\mathcal{Mon}$. □

5 Immediate consequences

The James Construction:

The underlying space functor $U : (\mathcal{Mon}, \mathcal{Mon}^w) \rightarrow (\mathcal{Top}^*, \mathcal{Top}^w)$ has a left adjoint

$$J : (\mathcal{Top}^*, \mathcal{Top}^w) \rightarrow (\mathcal{Mon}, \mathcal{Mon}^w)$$

commonly called the *James construction*, which associates with each based space X the free based monoid on X .

5.1 Proposition: (James [12]) For each path-connected based space there is a weak homotopy equivalence of spaces

$$JX \simeq \Omega\Sigma X.$$

D. Puppe investigated the conditions which would imply for this weak homotopy equivalence to be a genuine homotopy equivalence.

5.2 Proposition: (Puppe [7]): If X is a well-pointed path-connected Dold space then there is a homotopy equivalence.

$$JX \simeq \Omega\Sigma X.$$

Consider the diagram of functors

$$\begin{array}{ccc} \mathcal{Mon}^w & \begin{array}{c} \xleftarrow{B} \\ \xrightarrow{\Omega'} \end{array} & \mathcal{Top}^w \\ & \begin{array}{c} \searrow U \\ \swarrow J \end{array} & \begin{array}{c} \nearrow \Sigma \\ \nwarrow \Omega \end{array} \\ & \mathcal{Top}^w & \end{array}$$

All functors preserve weak equivalences. Hence they induce a diagram

$$\begin{array}{ccc} \text{Ho } \mathcal{Mon}^w & \begin{array}{c} \xleftarrow{\text{Ho } B} \\ \xrightarrow{\text{Ho } \Omega'} \end{array} & \text{Ho } \mathcal{Top}^w \\ & \begin{array}{c} \searrow \text{Ho } U \\ \swarrow \text{Ho } J \end{array} & \begin{array}{c} \nearrow \text{Ho } \Sigma \\ \nwarrow \text{Ho } \Omega \end{array} \\ & \text{Ho } \mathcal{Top}^w_i & \end{array}$$

consisting of adjoint pairs. Since the Moore loop space functor is naturally homotopy equivalent to the usual loop space functor there is a natural transformation

$$\tau(X) : U \circ \Omega'(X) \rightarrow \Omega(X)$$

which is a homotopy equivalence. Hence $\mathrm{Ho} \Omega$ and $\mathrm{Ho} U \circ \mathrm{Ho} \Omega'$ are naturally isomorphic. Since their left adjoints are unique up to natural isomorphisms this implies that $\mathrm{Ho} B \circ \mathrm{Ho} J$ and $\mathrm{Ho} \Sigma$ are naturally isomorphic. We obtain

5.3 Proposition: For each $X \in \mathcal{Top}^w$ there is a homotopy equivalence

$$BJ(X) \simeq \Sigma(X)$$

natural up to homotopy. $\square$

We obtain Puppe's result by combining 5.3 with another well-known result:

5.4 Proposition: If M is a well-pointed monoid whose underlying space is a Dold space and $\pi_0(M)$ is a group, then M is grouplike [7, (12.7)].

Proof of 5.2: If X is a path-connected Dold space, so is JX . Hence JX is grouplike and $\mu(WJX) : WJX \rightarrow W\Omega'BJX$ is a weak equivalence by 4.16, so that $\Omega'B\varepsilon(JX) \circ \varepsilon(\Omega'BJX) \circ \mu(WJX) \circ \iota(JX) : JX \rightarrow \Omega'BJX$ is a homotopy equivalence. We have a sequence of homotopy equivalences

$$JX \simeq \Omega'BJX \simeq \Omega BJX \simeq \Omega \Sigma X$$

Homotopical group completion: Homotopical group completion is the replacement of a monoid by a grouplike one having a universal property. We state our result for the full subcategory $\mathrm{Ho} \mathcal{Mon}^w$ of $\mathrm{Ho} \mathcal{Mon}$ of well-pointed monoids. Since $q(M) : VM \rightarrow M$ is a weak equivalence, $\mathrm{Ho} \mathcal{Mon}^w$ is equivalent to $\mathrm{Ho} \mathcal{Mon}$ so that the corresponding statement for $\mathrm{Ho} \mathcal{Mon}$ follows.

5.5 Proposition: Let M be a well-pointed monoid. The homotopy class of the homomorphism $\mu(WM) : WM \rightarrow W\Omega'BMW$, considered as a morphism in $\mathrm{Ho} \mathcal{Mon}^w(M, \Omega'BMW)$, is a group completion in the following sense: Given a diagram

$$\begin{array}{ccc} M & \xrightarrow{[\mu(WM)]} & \Omega'BMW \\ & \searrow [g] & \swarrow [\bar{g}] \\ & N & \end{array}$$

in $\mathrm{Ho} \mathcal{Mon}^w$ with N grouplike, there exists a unique morphism $[\bar{g}] : \Omega'BMW \rightarrow N$ making the diagram commute.

Proof Consider the homotopy commutative diagram in $\mathcal{Mon}^w$

$$\begin{array}{ccc} WM & \xrightarrow{\mu(WM)} & W\Omega'BMW \\ g \downarrow & & \downarrow W\Omega'Bg \\ WN & \xrightarrow{\mu(WN)} & W\Omega'BN \end{array}$$

Since N is well-pointed and grouplike $\mu(WN)$ is homotopy invertible in $\mathcal{Mon}^w$ by 4.16. We choose a homotopy inverse $h : W\Omega'BN \rightarrow WN$ and define $\bar{g} = h \circ W\Omega'Bg$. Then $\bar{g} \circ \mu(WM) \simeq g$ in $\mathcal{Mon}^w$.

For the uniqueness of $[\bar{g}]$ suppose there is a homomorphism $g' : W\Omega'BM \rightarrow WN$ such that $h \circ \mu(WM) \simeq g'$. Put $j_1 = \mu(WN) \circ g'$ and $j_2 = W\Omega'Bg$. It suffices to show that $j_1 \simeq j_2$ in $\mathcal{Mon}^w$. Since $Bj_1 \circ B\mu(WM) \simeq \mu(WN) \circ Bg \simeq Bj_2 \circ B\mu(WM)$ and $B\mu(WM)$ is a homotopy equivalence by 4.16, we obtain $Bj_1 \simeq Bj_2$. Since $\mu(W\Omega'BM)$ and $\mu(W\Omega'BN)$ are homotopy equivalences in $\mathcal{Mon}^w$ by 4.16 and μ is natural up to homotopy the following diagram is homotopy commutative and establishes the result:

$$\begin{array}{ccc} W\Omega'BM & \xrightarrow{\mu(W\Omega'BM)} & W\Omega'BW\Omega'BM \\ j_k \downarrow & & \downarrow W\Omega'Bj_k \\ W\Omega'BN & \xrightarrow{\mu(W\Omega'BN)} & W\Omega'BW\Omega'BN \end{array}$$

□

Dold spaces and grouplike monoids

For details on Dold spaces see [17]. We restrict our attention to the well-pointed case. Using the whiskering process it is easy to extend our results to the general case.

Let $\mathcal{Top}_{Dold}^w \subset \mathcal{Top}^w$ denote the full subcategory of well-pointed path-connected Dold spaces. Since BM is in $\mathcal{Top}_{Dold}^w$ for any well-pointed monoid by [17, Cor. 5.2], the classifying space functor restricts to a functor

$$B : \mathcal{Mon}^w \rightarrow \mathcal{Top}_{Dold}^w.$$

Let $\mathcal{Mon}_{group}^w \subset \mathcal{Mon}^w$ denote the full subcategory of grouplike well-pointed monoids. Then Proposition 4.16 implies

5.6 Theorem: The functors

$$B^{w\mathcal{H}} : \mathcal{HMon}_{group}^w \xleftrightarrow{\quad} \mathcal{HTop}_{Dold}^w : \Omega'^{w\mathcal{H}}$$

define an equivalence up to homotopy of categories, i.e. the natural transformations up to homotopy $\mu : \text{Id} \rightarrow \Omega'^{w\mathcal{H}} \circ B^{w\mathcal{H}}$ and $\eta : B^{w\mathcal{H}} \circ \Omega'^{w\mathcal{H}} \rightarrow \text{Id}$ take values in homotopy equivalences. In particular,

$$\text{Ho } B : \text{Ho } \mathcal{M}on_{group}^w \xleftrightarrow{\sim} \text{Ho } \mathcal{T}op_{Dold}^w : \text{Ho } \Omega'$$

define an equivalence of categories. $\square$

The second part is a slight extension of a well-known result (e.g. see [3, Section 4].

The following two propositions extend and strengthen results of Fuchs [8, Satz 7.7].

The diagram

$$\begin{array}{ccc} \mathcal{M}on^w(WM, WN) & \xrightarrow{\mu(WN)_*} & \mathcal{M}on^w(WM, W\Omega'BN) \\ & \searrow B & \downarrow \lambda(WM, BWN) \\ & & \mathcal{T}op^w(BWM, BWN) \end{array}$$

commutes up to homotopy because by 4.13

$$\begin{aligned} (\lambda(WM, BWN) \circ \mu(WN)_*)(f) &= \eta(BWN) \circ B' \circ \mu(WN)_*(f) \\ &= \eta(BWN) \circ B\mu(WN) \circ Bf \\ &\simeq Bf \quad \text{continuously in } f \end{aligned}$$

with $B' : \mathcal{M}on^w(WM, W\Omega'BN) \rightarrow \mathcal{T}op^w(BWM, BW\Omega'BN)$. If N is grouplike $\mu(WN)$ is a homotopy equivalence in $\mathcal{M}on^w$, and we obtain

5.7 Proposition: If N is a well-pointed grouplike monoid then

$$B : \mathcal{M}on^w(WM, WN) \rightarrow \mathcal{T}op^w(BWM, BWN)$$

is a homotopy equivalence. $\square$

Since $\eta(X) : BW\Omega'X \rightarrow X$ is a natural transformation the following diagram commutes

$$\begin{array}{ccc} \mathcal{M}on^w(W\Omega'X, W\Omega'Y) & \xleftarrow{W\Omega'} & \mathcal{T}op^w(X, Y) \\ \downarrow B & & \downarrow \eta(X)^* \\ \mathcal{T}op^w(BW\Omega'X, BW\Omega'Y) & \xrightarrow{\eta(Y)^*} & \mathcal{T}op^w(BW\Omega'X, Y) \end{array}$$

Since $W\Omega'Y$ is grouplike the map B is a homotopy equivalence by 5.7. Since $\eta(Y)_* \circ B = \lambda(W\Omega'X, Y)$ the map $\eta(Y)_* : \mathcal{Top}^w(BW\Omega'X, BW\Omega'Y) \rightarrow \mathcal{Top}^w(BW\Omega'X, Y)$ is a homotopy equivalence. If X is a well-pointed path-connected Dold space $\eta(X)$ is a based homotopy equivalence by 4.16. We obtain

5.8 Proposition: If X is a well-pointed path-connected Dold space then $W\Omega' : \mathcal{Top}^w(X, Y) \rightarrow \mathcal{Mon}^w(W\Omega'X, W\Omega'Y)$ is a homotopy equivalence. $\square$

Homotopy homomorphisms and unitary homotopy homomorphisms

5.9 Proposition: Let M and N be well-pointed monoids and N be group-like. Then $\varepsilon'(M) : \overline{W}M \rightarrow WM$ induces a homotopy equivalence

$$\mathcal{Mon}(WM, N) \rightarrow \mathcal{Sgp}(\overline{W}M, N).$$

Proof By 2.7 we may replace N by WN . Since $\mathcal{Sgp}(\overline{W}M, WN)$ is naturally homeomorphic to $\mathcal{Mon}(W(M_+), WN)$ by 2.5 it suffices to show that the counit $\kappa(M) : M_+ \rightarrow M$ induces a homotopy equivalence

$$\kappa(M)^* : \mathcal{Mon}(WM, WN) \rightarrow \mathcal{Mon}(W(M_+), WN)$$

The diagram

$$\begin{array}{ccc} \mathcal{Mon}(WM, WN) & \xrightarrow{B} & \mathcal{Top}^*(BWM, BWN) \\ \kappa(M)^* \downarrow & & \downarrow B\kappa(M)^* \\ \mathcal{Mon}(W(M_+), WN) & \xrightarrow{B} & \mathcal{Top}^*(BW(M_+), BWN) \end{array}$$

commutes. By 5.7 the maps B are homotopy equivalences, and by 4.2 the map $B\kappa(M)^*$ is a homotopy equivalence. Hence so is $\kappa(M)^*$. $\square$

5.10 Remark: In general we cannot expect that $\varepsilon'(M) : \overline{W}M \rightarrow WM$ induces a homotopy equivalence. E.g. it can happen that a homomorphism $\overline{W}M \rightarrow N$ does not map (e_M) into the path-component of e_N so that there is no chance to homotop it into a homomorphism $WM \rightarrow N$.

5.11 Proposition: If M is a well-pointed monoid then $Wq(M) : WVM \rightarrow WM$ is a homotopy equivalence in $\mathcal{Mon}^w$ by 2.12 inducing a homotopy equivalence

$$\mathcal{Mon}(WM, N) \rightarrow \mathcal{Mon}(WVM, N).$$

6 Diagrams of monoids

We want to show that the homotopy adjunction of Theorem 4.5 lifts to diagram categories. This is not evident: since the unit of our homotopy adjunction is only natural up to homotopy it does not lift to diagrams.

Let $\mathcal{M}$ be a cocomplete $\mathcal{T}op$ -enriched tensored category with a class $\mathcal{W}$ of weak equivalences containing the homotopy equivalences. We assume that $\mathcal{M}$ has a strong cofibrant replacement functor (Q_M, ε_M) . We use $\otimes$ for the tensor in $\mathcal{M}$ and Q for Q_M as long as there no ambiguity.

6.1 Definition: Let $\mathcal{C}$ be a small indexing category. A morphism $f : D_1 \rightarrow D_2$ of $\mathcal{C}$ -diagrams in $\mathcal{M}$ is called a *weak equivalence* if it is objectwise a weak equivalence in $\mathcal{M}$. We denote the class of weak equivalence in $\mathcal{M}^{\mathcal{C}}$ by $\mathcal{W}^{\mathcal{C}}$.

Our first aim is to show that $\mathcal{M}^{\mathcal{C}}$ admits a strong cofibrant replacement functor. We proceed as in Section 2.

We define a $\mathcal{C} \times \mathcal{C}^{\text{op}}$ -diagram $B(\mathcal{C}, \mathcal{C}, \mathcal{C})$ in $\mathcal{T}op$ as follows:

$$B(\mathcal{C}, \mathcal{C}, \mathcal{C})(b, a) = B(\mathcal{C}(-, b), \mathcal{C}, \mathcal{C}(a, -))$$

where the right side is the 2-sided bar construction of 4.1.

The $\mathcal{C} \times \mathcal{C}^{\text{op}}$ structure on $B_n(\mathcal{C}, \mathcal{C}, \mathcal{C})$ is given by

$$(g, h) \cdot (f_0, \dots, f_{n+1}) = (g \circ f_0, f_1, \dots, f_n, f_{n+1} \circ h)$$

Analogously we define a $\mathcal{C}^{\text{op}}$ -diagram $B(*, \mathcal{C}, \mathcal{C})$ in $\mathcal{T}op$, where $*$ denotes the constant $\mathcal{C}^{\text{op}}$ -diagram on a single point.

6.2 Lemma: Let X and Y be $\mathcal{C} \times \mathcal{C}^{\text{op}}$ -diagrams in $\mathcal{T}op$, let $p : X \rightarrow Y$ be a map of diagrams which is objectwise a homotopy equivalence. Then p induces a homotopy equivalence

$$p_* : \mathcal{T}op^{\mathcal{C} \times \mathcal{C}^{\text{op}}}(B(\mathcal{C}, \mathcal{C}, \mathcal{C}), X) \rightarrow \mathcal{T}op^{\mathcal{C} \times \mathcal{C}^{\text{op}}}(B(\mathcal{C}, \mathcal{C}, \mathcal{C}), Y)$$

in $\mathcal{T}op$.

Proof We apply the HELP-Lemma. So given a diagram

$$\begin{array}{ccc} K & \xrightarrow{\bar{f}_K} & \mathcal{T}op^{\mathcal{C} \times \mathcal{C}^{\text{op}}}(B(\mathcal{C}, \mathcal{C}, \mathcal{C}), X) \\ \downarrow i & & \downarrow p_* \\ L & \xrightarrow{\bar{g}} & \mathcal{T}op^{\mathcal{C} \times \mathcal{C}^{\text{op}}}(B(\mathcal{C}, \mathcal{C}, \mathcal{C}), Y) \end{array}$$

which commutes up to a homotopy $\bar{h}_{K,t} : \bar{g} \circ i \simeq p_* \circ \bar{f}_K$, where i is a closed cofibration, we have to construct extensions

$$\begin{aligned} \bar{f} : L &\rightarrow \mathcal{Top}^{\mathcal{C} \times \mathcal{C}^{\text{op}}}(B(\mathcal{C}, \mathcal{C}, \mathcal{C}), X) \\ \bar{h}_t : L &\rightarrow \mathcal{Top}^{\mathcal{C} \times \mathcal{C}^{\text{op}}}(B(\mathcal{C}, \mathcal{C}, \mathcal{C}), Y) \end{aligned}$$

of $\bar{f}_K$ respectively $\bar{h}_{K,t}$ such that $\bar{h}_t : \bar{g} \simeq p_* \circ \bar{f}$.

Taking adjoints the above diagram translates to the following diagram of $\mathcal{C} \times \mathcal{C}^{\text{op}}$ -spaces

$$\begin{array}{ccc} K \times B(\mathcal{C}, \mathcal{C}, \mathcal{C}) & \xrightarrow{f'} & X \\ i \times \text{id} \downarrow & & \downarrow p_* \\ L \times B(\mathcal{C}, \mathcal{C}, \mathcal{C}) & \xrightarrow{g} & Y \end{array}$$

which commutes up to a homotopy $h'_t : g \circ (i \times \text{id}) \simeq p_* \circ f'$ in $\mathcal{Top}^{\mathcal{C} \times \mathcal{C}^{\text{op}}}$, and it suffices to construct extensions $f : L \times B(\mathcal{C}, \mathcal{C}, \mathcal{C}) \rightarrow X$ of f' and $h_t : L \times B(\mathcal{C}, \mathcal{C}, \mathcal{C}) \rightarrow Y$ of h'_t such that $h_t : g \simeq p_* \circ f$ in $\mathcal{Top}^{\mathcal{C} \times \mathcal{C}^{\text{op}}}$.

We construct these extensions by induction on the natural filtration F_n of $L \times B(\mathcal{C}, \mathcal{C}, \mathcal{C})$ induced by the realization of the simplicial set $B_\bullet(\mathcal{C}, \mathcal{C}, \mathcal{C})$. We start with $F_0 = \coprod_{a,b,c} L \times \mathcal{C}(c, b) \times \mathcal{C}(a, c)$. The diagram

$$\begin{array}{ccc} K \times \{(\text{id}_c, \text{id}_c)\} & \xrightarrow{f'} & X(c, c) \\ \downarrow & & \downarrow p_* \\ L \times \{(\text{id}_c, \text{id}_c)\} & \xrightarrow{g} & Y(c, c) \end{array}$$

commutes up to a homotopy given by h'_t . Since $p_* : X(c, c) \rightarrow Y(c, c)$ is a homotopy equivalence and $K \rightarrow L$ is a closed cofibration the required extensions exist by the HELP-Lemma. We extend f over all of F_0 by $f(l, j_0, j_1) = X(j_0, j_1) \circ f(l, \text{id}, \text{id})$ and analogously for h_t .

Now suppose that f and h_t have been defined on F_{n-1} . We obtain F_n from F_{n-1} by attaching spaces $L \times (j_0, \dots, j_{n+1}) \times \Delta^n$ along $L \times (j_0, \dots, j_{n+1}) \times \partial \Delta^n$. Here the j_k are morphisms in $\mathcal{C}$ such that the composition

$$j_0 \circ \dots \circ j_{n+1} : a \rightarrow c_n \rightarrow \dots \rightarrow c_0 \rightarrow b$$

is defined and $j_1, \dots, j_n$ are not identities. Hence the extension f and the homotopy h_t are already defined on

$$\begin{aligned} D(L \times (j_0, j_1, \dots, j_n, j_{n+1}) \times \Delta^n) \\ = K \times (j_0, j_1, \dots, j_n, j_{n+1}) \times \Delta^n \cup L \times (j_0, j_1, \dots, j_n, j_{n+1}) \times \partial \Delta^n. \end{aligned}$$

We apply the HELP-Lemma to the homotopy commutative diagram

$$\begin{array}{ccc} D(L \times (\text{id}_{c_0}, j_1, \dots, j_n, \text{id}_{c_n}) \times \Delta^n) & \xrightarrow{f''} & X(c_0, c_n) \\ \downarrow & & \downarrow p_* \\ L \times (\text{id}_{c_0}, j_1, \dots, j_n, \text{id}_{c_n}) \times \Delta^n & \xrightarrow{g} & Y(c_0, c_n) \end{array}$$

where f'' and the commuting homotopy are given by the already defined extensions. Since p_* is a homotopy equivalence and the inclusion

$$D(L \times (\text{id}_{c_0}, j_1, \dots, j_n, \text{id}_{c_n}) \times \Delta^n) \subset L \times (\text{id}_{c_0}, j_1, \dots, j_n, \text{id}_{c_n}) \times \Delta^n$$

is a closed cofibration the required extensions exist. We extend our maps to maps of diagrams as in the F_0 -case. $\square$

Let D be a $\mathcal{C}$ -diagram in $\mathcal{M}$ and X a $\mathcal{C}^{\text{op}}$ -diagram in $\mathcal{Top}$. We define $X \otimes_{\mathcal{C}} D$ to be the coequalizer in $\mathcal{M}$ of

$$\coprod_{f \in \text{mor } \mathcal{C}} X(\text{target}(f)) \otimes D(\text{source}(f)) \xrightleftharpoons[\beta]{\alpha} \coprod_{a \in \text{ob } \mathcal{C}} X(a) \otimes D(a)$$

where for $f : a \rightarrow b$ in $\mathcal{C}$ the f -summand $X(b) \otimes D(a)$ is mapped as follows

$$\begin{aligned} \alpha &= X(f) \otimes \text{id} : X(b) \otimes D(a) \longrightarrow X(a) \otimes D(a) \\ \beta &= \text{id} \otimes D(f) : X(b) \otimes D(a) \longrightarrow X(b) \otimes D(b) \end{aligned}$$

We define a functor

$$R : \mathcal{M}^{\mathcal{C}} \rightarrow \mathcal{M}^{\mathcal{C}}, \quad D \mapsto B(\mathcal{C}, \mathcal{C}, \mathcal{C}) \otimes_{\mathcal{C}} QD$$

where $B(\mathcal{C}, \mathcal{C}, \mathcal{C}) \otimes_{\mathcal{C}} D$ is the $\mathcal{C}$ -diagram

$$a \mapsto B(\mathcal{C}(-, a), \mathcal{C}, \mathcal{C}) \otimes_{\mathcal{C}} D$$

in $\mathcal{M}$.

6.3 Proposition: Let D_0 , D_1 , and D_2 be $\mathcal{C}$ diagrams in $\mathcal{M}$, let $p : D_1 \rightarrow D_2$ be a weak equivalence in $\mathcal{M}^{\mathcal{C}}$ and $q : A_1 \rightarrow A_2$ a weak equivalence in $\mathcal{M}$. Then p and q induce homotopy equivalences

$$\begin{aligned} p_* : \mathcal{M}^{\mathcal{C}}(RD_0, D_1) &\rightarrow \mathcal{M}^{\mathcal{C}}(RD_0, D_2) \\ q_* : \mathcal{M}(B(*, \mathcal{C}, \mathcal{C}) \otimes_{\mathcal{C}} QD_0, A_1) &\rightarrow \mathcal{M}(B(*, \mathcal{C}, \mathcal{C}) \otimes_{\mathcal{C}} QD_0, A_2) \end{aligned}$$

in $\mathcal{Top}$.

Proof Since $\mathcal{M}^{\mathcal{C}}(RD_0, D_i) \cong \mathcal{Top}^{\mathcal{C} \times \mathcal{C}^{\text{op}}}(B(\mathcal{C}, \mathcal{C}, \mathcal{C}), \mathcal{M}(QD_0, D_i))$ it follows from Lemma 6.2 with $X(b, a) = \mathcal{M}(QD_0(a), D_1(b))$ and $Y(b, a) = \mathcal{M}(QD_0(a), D_2(b))$ that p_* is a homotopy equivalence.

There is a sequence of natural homeomorphisms

$$\begin{aligned} \mathcal{M}(B(*, \mathcal{C}, \mathcal{C}) \otimes_{\mathcal{C}} QD_0, A_i) &\cong \mathcal{Top}^{\mathcal{C}^{\text{op}}}(B(*, \mathcal{C}, \mathcal{C}), \mathcal{M}(QD_0, A_i)) \\ &\cong \mathcal{Top}^{\mathcal{C}^{\text{op}}}(\text{colim}_{\mathcal{C}} B(\mathcal{C}, \mathcal{C}, \mathcal{C}), \mathcal{M}(QD_0, A_i)) \\ &\cong \mathcal{Top}^{\mathcal{C} \times \mathcal{C}^{\text{op}}}(B(\mathcal{C}, \mathcal{C}, \mathcal{C}), \mathcal{M}(QD_0, \text{const} A_i)) \end{aligned}$$

where $\text{const} A_i$ are the constant $\mathcal{C}$ -diagrams on A_i . As in the first part, it follows that q_* is a homotopy equivalence. $\square$

Let $\mathcal{C}_{\bullet}$ denote the $\mathcal{C} \times \mathcal{C}^{\text{op}}$ -diagram of simplicial sets sending (a, b) to the constant simplicial set $\mathcal{C}(a, b)$. The maps

$$\begin{aligned} \delta_n : B_n(\mathcal{C}, \mathcal{C}, \mathcal{C})(a, b) &\longrightarrow \mathcal{C}(a, b) \\ (f_0, \dots, f_{n+1}) &\longmapsto f_0 \circ \dots \circ f_{n+1} \end{aligned}$$

define a simplicial map $B_{\bullet}(\mathcal{C}, \mathcal{C}, \mathcal{C}) \rightarrow \mathcal{C}_{\bullet}$. Let $\delta : B(\mathcal{C}, \mathcal{C}, \mathcal{C}) \rightarrow \mathcal{C}$ be its realization.

6.4 Proposition: $\delta(D) = \delta \otimes_{\mathcal{C}} \text{id}_D : B(\mathcal{C}, \mathcal{C}, \mathcal{C}) \otimes_{\mathcal{C}} D \rightarrow \mathcal{C} \otimes_{\mathcal{C}} D \cong D$ is objectwise a homotopy equivalence in $\mathcal{M}$ and hence a weak equivalence in $\mathcal{M}^{\mathcal{C}}$.

The proposition is an immediate consequence of the following Lemma:

6.5 Lemma: For each object $b \in \mathcal{C}$ the map $\varepsilon : B(\mathcal{C}, \mathcal{C}, \mathcal{C})(-, b) \rightarrow \mathcal{C}(-, b)$ is a homotopy equivalence in the category $\mathcal{Top}^{\mathcal{C}^{\text{op}}}$.

Proof For $a \in \mathcal{C}$ let $\mathcal{X}_a$ denote the category whose objects are diagrams $a \xrightarrow{j_1} c \xrightarrow{j_0} b$ and whose morphisms from this object to $a \xrightarrow{j'_1} c' \xrightarrow{j'_0} b$ are morphisms $h : c \rightarrow c'$ in $\mathcal{C}$ making the diagram

$$\begin{array}{ccccc} & & c & & \\ & \nearrow^{j_1} & \downarrow h & \searrow^{j_0} & \\ a & & & & b \\ & \searrow_{j'_1} & & \nearrow_{j'_0} & \\ & & c' & & \end{array}$$

commute. Let $\mathcal{C}(a, b)$ stand for the discrete category whose object set is $\mathcal{C}(a, b)$. Then

$$\varepsilon_a : \mathcal{X}_a \rightarrow \mathcal{C}(a, b), \quad (a \xrightarrow{j_1} c \xrightarrow{j_0} b) \mapsto (j_0 \circ j_1 : a \rightarrow b)$$

defines a functor which has the section

$$s_a : \mathcal{C}(a, b) \rightarrow \mathcal{X}_a, \quad j \mapsto (a \xrightarrow{j} b \xrightarrow{\text{id}} b).$$

There is a natural transformation $\tau_a : \text{Id}_{\mathcal{X}_a} \rightarrow s_a \circ \varepsilon_a$ defined by the diagram

$$\begin{array}{ccccc} & & c & & \\ & \nearrow j_1 & \downarrow j_0 & \nwarrow j_0 & \\ a & & & & b \\ & \searrow j_0 \circ j_1 & & \nearrow \text{id} & \\ & & b & & \end{array}$$

So ε_a induces a homotopy equivalence of the classifying spaces. Now $B(\mathcal{X}_a) = B(\mathcal{C}, \mathcal{C}, \mathcal{C})(a, b)$ and $B(\mathcal{C}(a, b) = \mathcal{C}(a, b)$. Moreover all data are natural with respect to $a \in \mathcal{C}^{\text{op}}$. Hence we obtain the required result. $\square$

6.6 Corollary: $R : \mathcal{M}^c \rightarrow \mathcal{M}^c$ together with $\epsilon = \delta \otimes_{\mathcal{C}} \varepsilon_M : R \rightarrow \text{Id}$ is a strong cofibrant replacement functors.

Let $\mathcal{N}$ be another cocomplete $\mathcal{T}op$ -enriched tensored category with a class of weak equivalences containing the homotopy equivalences and a strong cofibrant replacement functor (Q_N, ε_N) .

6.7 Theorem: Let

$$F : \mathcal{M} \rightleftarrows \mathcal{N} : G$$

be continuous functors inducing a natural homotopy equivalence

$$\lambda(Q_M A, Q_N X) : \mathcal{M}(Q_M A, Q_M G Q_N X) \rightarrow \mathcal{N}(Q_N F Q_M A, Q_N X)$$

so that

$$F^{\mathcal{H}} : \mathcal{H}\mathcal{M} \rightleftarrows \mathcal{H}\mathcal{N} : G^{\mathcal{H}}$$

is a conatural adjunction up to homotopy. Then

$$(F^c)^{\mathcal{H}} : \mathcal{H}(\mathcal{M}^c) \rightleftarrows \mathcal{H}(\mathcal{N}^c) : (G^c)^{\mathcal{H}}$$

is an adjunction up to homotopy, and hence

$$\text{Ho}(F^c) : \text{Ho}(\mathcal{M}^c) \rightleftarrows \text{Ho}(\mathcal{N}^c) : \text{Ho}(G^c)$$

a genuine adjunction.

Proof For diagrams $D : \mathcal{C} \rightarrow \mathcal{M}$ and $Z : \mathcal{C} \rightarrow \mathcal{N}$ we have a sequence of natural maps

$$\begin{array}{ccc}
\mathcal{Top}^{\mathcal{C} \times \mathcal{C}^{\text{op}}} (B(\mathcal{C}, \mathcal{C}, \mathcal{C}), \mathcal{N}(Q_N F Q_M D, Q_N Z)) & \xrightarrow{\cong} & \mathcal{N}^{\mathcal{C}}(R_N F Q_M D, Q_N Z) \\
\uparrow \lambda(Q_M D, Q_N Z)_* & & \downarrow R_N F \delta(Q_M D)_* \\
\mathcal{Top}^{\mathcal{C} \times \mathcal{C}^{\text{op}}} (B(\mathcal{C}, \mathcal{C}, \mathcal{C}), \mathcal{M}(Q_M D, Q_M G Q_N Z)) & & \mathcal{N}^{\mathcal{C}}(R_N F R_M D, Q_N Z) \\
\uparrow \cong & & \uparrow \delta(Q_N Z)_* \\
\mathcal{M}^{\mathcal{C}}(R_M D, Q_M G Q_N Z) & & \mathcal{N}^{\mathcal{C}}(R_N F R_M D, R_N Z) \\
\uparrow \delta(Q_M G Q_N Z)_* & & \\
\mathcal{M}^{\mathcal{C}}(R_M D, R_M G Q_N Z) & & \\
\uparrow R_M G \delta(Q_N Z)_* & & \\
\mathcal{M}^{\mathcal{C}}(R_M D, R_M G R_N Z) & &
\end{array}$$

By assumption $\lambda(Q_M D, Q_N Z)$ is a homotopy equivalence. Since $\delta(D)$ is objectwise a homotopy equivalence and since continuous functors preserve homotopy equivalences, $R_M G \delta(Q_N Z)$ and $R_N F \delta(Q_M D)$ are homotopy equivalences in $\mathcal{M}^{\mathcal{C}}$ by 6.3 so that $R_M G \delta(Q_N Z)_*$ and $R_N F \delta(Q_M D)_*$ are homotopy equivalences in $\mathcal{Top}$, and $\delta(Q_N Z)_*$ and $\delta(Q_M G Q_N Z)_*$ are homotopy equivalences in $\mathcal{Top}$ by 6.3. $\square$

6.8 Addendum: The last natural map in the proof of the theorem points in the wrong direction. So we cannot conclude that $(F^{\mathcal{C}})^{\mathcal{H}}$ and $(G^{\mathcal{C}})^{\mathcal{H}}$ are a conatural homotopy adjoint pair.

$$\eta(Q_N Y) = \lambda(Q_M G Q_N Y, Q_N Y)(\text{id}_{Q_M G Q_N Y}) : Q_N F Q_M G Q_N Y \rightarrow Q_N Y$$

is natural with respect to morphisms $f : Q_N Y_1 \rightarrow Q_N Y_2$ in $\mathcal{N}$. If η extends to a natural map $\eta(Y) : Q_N F Q_M G Y \rightarrow Y$ for all $Y \in \mathcal{N}$ or at least for all Y of the form $Y = R_N Y'$ we obtain a natural map $\lambda^{\mathcal{C}}(R_M D, R_N Z)$ defined by

$$\begin{array}{ccc}
\mathcal{M}^{\mathcal{C}}(R_M D, R_M G R_N Z) & \xrightarrow{R_N F} & \mathcal{M}^{\mathcal{C}}(R_N F R_M D, R_N F R_M G R_N Z) \\
& \searrow \lambda^{\mathcal{C}}(R_M D, R_N Z) & \downarrow \eta(R_N Z) \\
& & \mathcal{N}^{\mathcal{C}}(R_N F R_M D, R_N Z)
\end{array}$$

which makes the diagram of the proof of the theorem commute so that $(F^{\mathcal{C}})^{\mathcal{H}}$ and $(G^{\mathcal{C}})^{\mathcal{H}}$ are a conatural homotopy adjoint pair.

For use in the next proposition we note

6.9 Lemma: Let $D : \mathcal{C} \rightarrow \mathcal{Mon}^w$ is a diagram of well-pointed monoids. Then $B(*, \mathcal{C}, \mathcal{C}) \otimes_{\mathcal{C}} D$ is a well-pointed space, and $B(\mathcal{C}, \mathcal{C}, \mathcal{C}) \otimes_{\mathcal{C}} D$ and $B(\mathcal{C}, \mathcal{C}, \mathcal{C}) \otimes_{\mathcal{C}} WD$ are diagrams of well-pointed monoids.

Proof The first part holds by [15, Prop. 7.8]. The second and third statement follow by the argument used in [15, Prop. 7.8]. $\square$

From 6.6 and 6.9 we obtain

6.10 Proposition: With the choices of weak equivalences $\mathcal{W}$ as in 3.7 the functors

$$\begin{array}{lll} \mathcal{Mon}^{\mathcal{C}} & \rightarrow & \mathcal{Mon}^{\mathcal{C}} & D \mapsto B(\mathcal{C}, \mathcal{C}, \mathcal{C}) \otimes_{\mathcal{C}} WVD \\ (\mathcal{Mon}^w)^{\mathcal{C}} & \rightarrow & (\mathcal{Mon}^w)^{\mathcal{C}} & D \mapsto B(\mathcal{C}, \mathcal{C}, \mathcal{C}) \otimes_{\mathcal{C}} WD \\ \mathcal{Sgp}^{\mathcal{C}} & \rightarrow & \mathcal{Sgp}^{\mathcal{C}} & D \mapsto B(\mathcal{C}, \mathcal{C}, \mathcal{C}) \otimes_{\mathcal{C}} \overline{WD} \\ (\mathcal{Top}^*)^{\mathcal{C}} & \rightarrow & (\mathcal{Top}^*)^{\mathcal{C}} & D \mapsto B(\mathcal{C}, \mathcal{C}, \mathcal{C})_+ \wedge_{\mathcal{C}} V^t D \\ (\mathcal{Top}^w)^{\mathcal{C}} & \rightarrow & (\mathcal{Top}^w)^{\mathcal{C}} & D \mapsto B(\mathcal{C}, \mathcal{C}, \mathcal{C})_+ \wedge_{\mathcal{C}} D \end{array}$$

together with the corresponding natural transformations ϵ are strong replacement functors with respect to the weak equivalences in $\mathcal{W}^{\mathcal{C}}$. In particular, the localizations of these categories with respect to $\mathcal{W}^{\mathcal{C}}$ exist. (Recall that $K_+ \wedge X$ is the tensor over $\mathcal{Top}$ in $\mathcal{Top}^*$.)

Since Addendum 6.8 applies to our situation in Section 4 we obtain

6.11 Theorem: The homotopy adjunctions of Theorems 4.4 and 4.5 lift to conatural homotopy adjunctions

$$(B^{\mathcal{C}})^{\mathcal{H}} : \mathcal{H}(\mathcal{Mon}^w)^{\mathcal{C}} \rightleftarrows (\mathcal{Top}^w)^{\mathcal{C}} : (\Omega^{\mathcal{C}})^{\mathcal{H}}$$

and

$$(B^{\mathcal{C}})^{\mathcal{H}} : \mathcal{H}\mathcal{Mon}^{\mathcal{C}} \rightleftarrows \mathcal{Top}^{*\mathcal{C}} : (\Omega^{\mathcal{C}})^{\mathcal{H}}.$$

There are natural adjunction homotopy equivalences

$$\begin{array}{ll} \lambda(RD, QZ) : (\mathcal{Mon}^w)^{\mathcal{C}}(RD, R\Omega'QZ) & \rightarrow (\mathcal{Top}^w)^{\mathcal{C}}(QBRD, QZ) \\ \lambda(RVD, QV^tZ) : \mathcal{Mon}^{\mathcal{C}}(RVD, RV\Omega'QV^tZ) & \rightarrow \mathcal{Top}^{*\mathcal{C}}(QV^tBRVD, QV^tZ) \end{array}$$

in $\mathcal{Top}$, where (R, ϵ) and (Q, ϵ^t) are the cofibrant replacement functors in $(\mathcal{Mon}^w)^{\mathcal{C}}$ respectively $(\mathcal{Top}^w)^{\mathcal{C}}$ of 6.10. Hence

$$\mathrm{Ho} B^{\mathcal{C}} : \mathrm{Ho}(\mathcal{Mon}^w)^{\mathcal{C}} \rightleftarrows \mathrm{Ho}(\mathcal{Top}^w)^{\mathcal{C}} : \mathrm{Ho} \Omega^{\mathcal{C}}$$

and

$$\mathrm{Ho} B^{\mathcal{C}} : \mathrm{Ho}(\mathcal{M}on)^{\mathcal{C}} \rightleftarrows \mathrm{Ho}(\mathcal{T}op^*)^{\mathcal{C}} : \mathrm{Ho} \Omega'^{\mathcal{C}}$$

are genuine adjunctions.

6.12 Theorem: Let $\mathcal{M}$ be as above. Then the adjoint pair of functors

$$\mathrm{colim} : \mathcal{M}^{\mathcal{C}} \rightleftarrows \mathcal{M} : \mathrm{const}$$

induces a conatural adjunction up to homotopy

$$\mathrm{colim}^{\mathcal{H}} : \mathcal{H}\mathcal{M}^{\mathcal{C}} \rightleftarrows \mathcal{H}\mathcal{M} : \mathrm{const}^{\mathcal{H}}.$$

Hence we obtain a genuine adjunction

$$\mathrm{Ho} \mathrm{colim} : \mathrm{Ho} \mathcal{M}^{\mathcal{C}} \rightleftarrows \mathrm{Ho} \mathcal{M} : \mathrm{Ho} \mathrm{const}$$

Proof We have the following sequence of natural homotopy equivalences and homeomorphisms from $\mathcal{H}\mathcal{M}^{\mathcal{C}}(D, \mathrm{const}^{\mathcal{H}} A) = \mathcal{M}^{\mathcal{C}}(RD, R(\mathrm{const} QA))$ to $\mathcal{H}\mathcal{M}(\mathrm{colim}^{\mathcal{H}} D, A) = \mathcal{M}(Q(\mathrm{colim} RD), QA)$:

$$\begin{array}{lll} (1) & \mathcal{M}^{\mathcal{C}}(RD, R(\mathrm{const} QA)) & \xrightarrow{\epsilon(\mathrm{const} QA)_*} \mathcal{M}^{\mathcal{C}}(RD, \mathrm{const} QA) \\ (2) & & \xrightarrow{\cong} \mathcal{M}(\mathrm{colim} RD, QA) \\ (3) & & \xrightarrow{\epsilon_M(\mathrm{colim} RD)^*} \mathcal{M}(Q(\mathrm{colim} RD), QA). \end{array}$$

The first map is a homotopy equivalence by 6.3, the second one is the adjunction homeomorphism, and the third one is a homotopy equivalence, because $\epsilon_M(\mathrm{colim} RD) : Q \mathrm{colim} RD \rightarrow \mathrm{colim} RD$ is a homotopy equivalence in $\mathcal{M}$ by 6.3. $\square$

6.13 Definition: The *homotopy colimit functor* $\mathrm{hocolim} : \mathcal{M}^{\mathcal{C}} \rightarrow \mathcal{M}$ is defined by

$$\mathrm{hocolim} D = \mathrm{colim} RD = B(*, \mathcal{C}, \mathcal{C}) \otimes_{\mathcal{C}} QD.$$

6.14 Remark: In the literature one often finds the homotopy colimit defined by $\mathrm{hocolim} D = B(*, \mathcal{C}, \mathcal{C}) \otimes_{\mathcal{C}} D$ (e.g. see [10, 18.1.1]). This has historical reasons because homotopy colimits were first defined in categories where all object were cofibrant.

We apply these results to $\mathcal{M}on$ and prove

6.15 Theorem: The classifying space functor

$$B : (\mathcal{M}on, \mathcal{M}on^w) \rightarrow (\mathcal{T}op^*, \mathcal{T}op^w)$$

preserves homotopy colimits up to genuine homotopy equivalences. More precisely, for any diagram $D : \mathcal{C} \rightarrow \mathcal{M}on$ the natural map

$$\mathrm{hocolim}^{\mathcal{T}op^*} BD \rightarrow B(\mathrm{hocolim}^{\mathcal{M}on} D)$$

is a homotopy equivalence.

Proof By definition of the homotopy colimit functor it suffices to prove the well-pointed case.

Consider the diagram

$$\begin{array}{ccccc} (\mathcal{M}on^w)^{\mathcal{C}} & \xrightarrow{\gamma_{(\mathcal{M}on^w)^{\mathcal{C}}}} & \mathrm{Ho}(\mathcal{M}on^w)^{\mathcal{C}} & \xrightarrow{\mathrm{Ho\,colim}} & \mathrm{Ho}\,\mathcal{M}on \\ B^{\mathcal{C}} \downarrow & & \mathrm{Ho}\,B^{\mathcal{C}} \downarrow & & \downarrow \mathrm{Ho}\,B \\ (\mathcal{T}op^w)^{\mathcal{C}} & \xrightarrow{\gamma_{\mathcal{T}op^w}} & \mathrm{Ho}(\mathcal{T}op^*)^{\mathcal{C}} & \xrightarrow{\mathrm{Ho\,colim}} & \mathrm{Ho}\,\mathcal{T}op^w \end{array}$$

and recall that $\mathrm{Ho\,colim}$ is induced by the homotopy colimit functor. Since B preserves weak equivalences in the well-pointed case, $B^{\mathcal{C}}$ induces $\mathrm{Ho}\,B^{\mathcal{C}}$ so that the left square commutes up to natural equivalence. The right square commutes up to natural equivalence, because the corresponding square of right adjoints commutes. Hence, for any diagram D in $\mathcal{M}on$, the natural map

$$\mathrm{hocolim}^{\mathcal{T}op^w} BD \rightarrow B(\mathrm{hocolim}^{\mathcal{M}on^w} D)$$

becomes an isomorphism in $\mathrm{Ho}\,\mathcal{T}op^w = \pi\mathcal{T}op^w$. $\square$

References

- [1] G. Allaud, De-looping homotopy equivalences, Arch. Math. 23 (1972), 167 - 169.
- [2] C. Berger, I. Moerdijk, The Boardman-Vogt resolution of operads in monoidal model categories, Topology 45 (2006), 807-849.
- [3] M. Brinkmeier, Strongly homotopy-commutative monoids revisited, Documenta Mathematica 5 (2000), 613-624.

- [4] J.M. Boardman, R.M. Vogt, Homotopy invariant algebraic structures on topological spaces, Lecture Notes in Mathematics 347, Springer, 1973.
- [5] J. M. Boardman and R. M. Vogt, Tensor products of theories, applications to infinite loop spaces, J. Pure Appl. Algebra 14 (1979), 117-129.
- [6] F. Borceux, Handbook of categorical algebra 2, Encyclopedia of Mathematics and its Applications 51, Cambridge University Press 1994.
- [7] T. tom Dieck, K.H. Kamps, D. Puppe, Homotopietheorie, Springer Lecture Notes in Mathematics 157 (1970).
- [8] M. Fuchs, Verallgemeinerte Homotopie-Homomorphismen und klassifizierende Räume, Math. Annalen 161 (1965), 197-230.
- [9] M. Fuchs, Homotopy equivalences in equivariant topology, Proc. Amer. Math. Soc. 58 (1976), 347-352.
- [10] P.S. Hirschhorn, Model categories and their localizations, Mathematical Surveys and Monographs 99, Amer. Math. Soc. (2002).
- [11] P.G. Goerss, J.F. Jardine, Simplicial homotopy theory, Progress in Mathematics 174, Birkhäuser Verlag, Basel (1999).
- [12] I.M. James, Reduced product spaces, Ann. of Math. 62 (1955), 170-197.
- [13] G.M. Kelly, Basic concepts of enriched category theory, London Math. Soc. Lecture Note Series 64, Cambridge University Press (1982).
- [14] R.J. Milgram, The bar construction and abelian H -spaces, Illinois J. Math. 11 (1967), 242-250.
- [15] T. Panov, N. Ray, R.M. Vogt, Colimits, Stanley-Reisner algebras, and loop spaces, Progress in Math. 215 (2003), 261-291.
- [16] D. Puppe, Some well known weak homotopy equivalences are genuine homotopy equivalences, Symposia Mathematica 5, Istituto Nazionale di Alta Matematica (1971), 363 - 374.

- [17] E. Schwamberger, R.M. Vogt, Dold spaces in homotopy theory, *Algebraic & Geometric Topology* (2009), 1585-1622.
- [18] G. Segal, Categories and cohomology theories, *Topology* 13 (1974), 293-312.
- [19] A. Strøm, Note on cofibrations II, *Math. Scand.* 22 (1968), 130-142.
- [20] A. Strøm, The homotopy category is a homotopy category, *Arch. Math.* 23 (1972), 435-441.
- [21] M. Sugawara, On the homotopy-commutativity of groups and loop spaces, *Mem. Coll. Sci. Univ. Kyoto Ser. A.* 33 (1960), 257-269.
- [22] R.M. Vogt, Cofibrant operads and universal E_∞ -operads, *Top. Appl.* 133 (2003), 69-87.
- [23] R.M. Vogt, The HELP-Lemma and its converse in Quillen model categories, *J. Homotopy and Related Structures* 6 (2011), 115-118.